Journal of Cardiovascular Disease Research

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Cardiovascular Vulnerability to Air Pollution in India

Lalatendu Mohanty

Department of Pharmaceutical Sciences,
HNB Garhwal University, Uttarakhand.

Abstract

Background: India faces a dual burden of rapidly increasing cardiovascular disease (CVD) prevalence and severe air pollution, particularly in urban areas. Hence, understanding the relationship between these two is essential for developing effective prevention methods to address public health challenges.

Objectives: This review synthesizes available evidence on cardiovascular vulnerability to air pollution in India, examining exposure patterns, epidemiological associations, biological mechanisms, and regional disparities to inform policy and research priorities.

Methods: A comprehensive review of peer-reviewed literature was conducted. The main focus of the study was on air quality monitoring data from Indian cities, epidemiological studies linking air pollution to cardiovascular outcomes in India, and mechanistic studies elucidating the biological pathways underlying particulate matter-induced cardiovascular injury.

Results: Urban India experiences severe particulate air pollution, with PM10 concentrations frequently exceeding 200 ug/m3 in major cities. Limited epidemiological evidence from Delhi, Mumbai, and other urban areas suggests associations between air pollution and cardiovascular morbidity and mortality. Other pathways include oxidative stress, systemic inflammation, autonomic nervous system dysfunction, and enhanced coagulation and thrombosis also related to air pollution. Substantial regional disparities exist, with northern cities experiencing higher pollution levels than southern regions. Vulnerable populations include the elderly, individuals with pre-existing cardiovascular conditions, diabetics, and those of lower socioeconomic status.

Conclusions: Air pollution emerged as a cardiovascular risk factor in India. Urgent action is needed to strengthen air quality monitoring networks, conduct rigorous epidemiological studies in diverse Indian populations, elucidate population-specific susceptibilities, and implement evidence-based pollution control policies to protect cardiovascular health.

Keywords: *Air pollution; cardiovascular disease; particulate matter; India; oxidative stress; inflammation; regional disparities; PM10; PM2.5; urban health*

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Table of Contents

1. Introduction
2. Air Pollution in India
    2.1 Exposure Burden and Sources
    2.2 Monitoring Infrastructure and Data Availability
    2.3 PM10 and PM2.5 Trends in Urban India
3. Cardiovascular Disease Epidemiology in India
4. Evidence Linking Air Pollution to Cardiovascular Disease in India
    4.1 Epidemiological Studies from Delhi
    4.2 Evidence from Mumbai and Other Cities
    4.3 Limitations of Current Evidence
5. Biological Pathways Linking Air Pollution to Cardiovascular Disease
    5.1 Oxidative Stress
    5.2 Systemic Inflammation
    5.3 Autonomic Nervous System Dysfunction
    5.4 Coagulation and Thrombosis
6. Regional Disparities in Air Pollution and Cardiovascular Risk
    6.1 Delhi and North India
    6.2 Kolkata and East India
    6.3 Mumbai and West India
    6.4 South India
    6.5 Rural versus Urban Disparities
7. Vulnerable Populations
8. Policy Implications and Gaps
9. Conclusions
10. References

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

1. Introduction

Cardiovascular disease (CVD) has emerged as the leading cause of mortality in India, estimated around 32% of all deaths and representing a major shift in the country's disease burden from infectious to non-communicable diseases [25]. India also faces severe air pollution, particularly in rapidly expanding urban areas, where industrialisation, vehicular traffic, and biomass combustion contribute to hazardous levels of airborne particulate matter [5]. The convergence of these two public health crises—rising CVD prevalence and deteriorating air quality—presents a critical challenge for India's health system and economic development.

Globally, epidemiological evidence has established that air pollution, particularly fine particulate matter (PM2.5) and coarse particles (PM10), is an independent risk factor for cardiovascular morbidity and mortality [6,9,13]. Studies from North America and Europe have demonstrated associations between both short-term and long-term particulate exposure and increased risks of myocardial infarction, stroke, heart failure exacerbations, arrhythmias, and cardiovascular death [14,15,36]. Major health organisations, including the American Heart Association, now recognise air pollution as a modifiable cardiovascular risk factor that needs clinical and public health attention.

However, the vast majority of research on air pollution and cardiovascular health has been conducted in developed countries with relatively lower pollution levels and different population characteristics than those found in India. India's unique context—characterised by extreme pollution levels, distinct sources of particulate matter, high prevalence of traditional cardiovascular risk factors, genetic susceptibilities, and limited healthcare infrastructure—necessitates India-specific research to understand cardiovascular vulnerability to air pollution in this population [5,8].

Despite the high magnitude of the problem, research on air pollution and cardiovascular health in India remains limited. A small number of epidemiological studies have examined associations between air pollution and health outcomes in Indian cities, primarily focusing on respiratory effects [2,4,5]. However, the Cardiovascular endpoints have received far less attention, with only a handful of studies specifically investigating air pollution-CVD relationships in Indian populations [1,2,3]. This lack of evidence represents a knowledge gap that limits the development of evidence-based policies for cardiovascular health in India.

This review aims to synthesize the available evidence on cardiovascular vulnerability to air pollution in India by: (1) characterizing the exposure burden and trends in ambient particulate pollution across Indian regions; (2) reviewing epidemiological evidence linking air pollution to cardiovascular outcomes in Indian populations; (3) examining biological pathways through which particulate matter induces cardiovascular injury; (4) analyzing regional disparities in pollution levels and potential cardiovascular impacts; (5) identifying vulnerable populations at heightened risk; and (6) discussing policy implications and research priorities. By integrating exposure assessment, epidemiology, and mechanistic understanding, this review provides a comprehensive framework for understanding and addressing cardiovascular vulnerability to air pollution in India.

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

2. Air Pollution in India

2.1 Exposure Burden and Sources

India experiences some of the highest levels of air pollution globally, with particulate matter posing the greatest health concern [5,8,19]. The sources of particulate air pollution in India are diverse and complex. The country's rapid industrialisation, urbanisation, and continued reliance on traditional biomass fuels are contributor of Air pollution. Major sources include vehicular emissions, industrial activities, power generation, construction dust, biomass burning for cooking and heating, agricultural waste burning, and natural sources such as windblown dust [5].

Vehicular emissions are a primary source of urban air pollution in Indian cities, as vehicle numbers have increased rapidly in recent decades [5]. Unlike developed countries, where emission standards and fuel quality have improved substantially, many vehicles in India operate with older technologies and poor-quality fuels, resulting in high emissions of particulate matter, nitrogen oxides, carbon monoxide, and volatile organic compounds. Two-stroke engines, diesel vehicles, and poorly maintained vehicles are also responsible for increasing urban air pollution [5].

Industrial sources vary by region but include thermal power plants, cement factories, brick kilns, chemical industries, and small-scale manufacturing units [5]. Many industrial facilities in India lack adequate pollution control technologies, and enforcement of emission standards remains inconsistent. Biomass combustion for domestic cooking and heating, while primarily an indoor air pollution issue, also contributes to particulate levels, particularly in semi-urban and rural areas [22,29].

Seasonal factors also significantly influence air pollution levels in India. The post-monsoon and winter months typically experience the highest pollution levels due to meteorological conditions that limit dispersion, increased biomass burning, and festival-related activities such as Diwali fireworks [26]. Agricultural burning of crop residues, particularly in northern India, contributes to severe episodic pollution events [5].

The composition of particulate matter in Indian cities differs from that in developed countries, with higher proportions of crustal material, black carbon, and organic compounds from biomass combustion [5]. This compositional difference may have important implications for health effects, as toxicity varies with particle composition and size distribution. [8,32].

2.2 Monitoring Infrastructure and Data Availability

Air quality monitoring infrastructure in India has historically been limited, with significant gaps in spatial coverage, temporal continuity, and measurement of fine particulate matter (PM2.5) [5]. The National Air Quality Monitoring Programme (NAMP), established by the Central Pollution Control Board (CPCB), serves as the primary systematic monitoring effort, but the network's coverage remains inadequate given India's population and geographic diversity [5].

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Most monitoring stations in India measure PM10 (particulate matter with aerodynamic diameter <=10 um) and total suspended particulates (TSP), with limited measurement of PM2.5, which is more strongly associated with cardiovascular health effects [5,8]. The monitoring network is concentrated in major urban centres, with rural and semi-urban areas largely unmonitored. Furthermore, many monitoring stations experience operational challenges, including equipment malfunctions, irregular maintenance, and gaps in data collection, limiting the availability of continuous, high-quality data for epidemiological research [5].

The lack of comprehensive monitoring data poses significant challenges for exposure assessment in epidemiological studies and for evaluating the effectiveness of pollution control interventions. Most available data come from regulatory monitoring stations that may not adequately capture population exposure, particularly in areas with high spatial variability in pollution levels [5]. Personal exposure monitoring and source apportionment studies remain rare in India, further limiting understanding of exposure patterns and health-relevant pollution sources.

2.3 PM10 and PM2.5 Trends in Urban India

Available monitoring data reveal alarmingly high levels of particulate air pollution in Indian cities, with concentrations frequently exceeding national and international air quality standards [5,8]. Annual average PM10 concentrations in major Indian cities typically range from 100 to 400 µg/m3, far exceeding the Indian National Air Quality Standard of 60 µg/m3 and the World Health Organisation guideline of 20 µg/m3 [5]. Delhi, India's capital and one of the world's most polluted megacities, consistently records some of the highest PM10 levels globally. Studies from the late 1990s and early 2000s reported annual average PM10 concentrations ranging from 150 to 300 ug/m3, with peak daily values exceeding 500 ug/m3 during winter months [2,4,5]. These extreme levels are due to the combined influence of vehicular emissions, industrial sources, construction activities, and unfavourable meteorological conditions.

Mumbai, India's financial capital, also experiences severe particulate pollution, though levels are generally lower than in Delhi. Studies have reported annual average PM10 concentrations ranging from 100 to 200 ug/m3 in different areas of the city, with higher levels in industrial and traffic-congested areas [3]. A case study in Mumbai documented significant health impacts associated with PM10 exposure. That highlighted the public health burden of particulate pollution in this megacity [3].

Kolkata, in eastern India, faces substantial particulate pollution from vehicular traffic, industrial emissions, and domestic biomass burning. Monitoring studies have documented high PM10 and TSP levels, with significant seasonal variation and differences between residential and industrial areas [5]. The city's dense population and limited green space exacerbate exposure to air pollution.

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Limited data on PM2.5 trends in Indian cities suggest that fine particles constitute a substantial fraction of PM10, typically 40-60%, though this varies by location and season [5]. The high proportion of fine particles is particularly concerning from a cardiovascular health perspective, as PM2.5 penetrates deeper into the respiratory system and has greater potential for systemic effects [7,10,11].

Temporal trends in particulate pollution in Indian cities are difficult to assess due to limited long-term monitoring data. However, available evidence suggests that pollution levels have remained persistently high or increased in many cities over the past two decades, driven by continued growth in vehicular traffic, industrial activity, and urban population [5].

3. Cardiovascular Disease Epidemiology in India

India is experiencing a dramatic increase in cardiovascular disease burden. CVD has emerged as the leading cause of death in India, with ischemic heart disease and stroke accounting for a substantial proportion of mortality and disability [25]. Demographic changes, urbanisation, lifestyle modifications, and shifts in traditional risk factors contribute to this.

The prevalence of coronary heart disease (CHD) in India has increased substantially over recent decades, with urban areas experiencing higher rates than rural regions [25]. Estimates suggest that CHD prevalence ranges from 3% to 4% in rural areas to 8% to 10% in urban areas.[25]. Stroke prevalence similarly shows an urban-rural gradient, with higher rates in cities.

Traditional cardiovascular risk factors—including hypertension, diabetes mellitus, dyslipidemia, obesity, tobacco use, and physical inactivity—have increased in prevalence across India, particularly in urban populations [25]. These conditions represent established risk factors for CVD and may also modify susceptibility to cardiovascular effects induced by air pollution [39,47].

The age distribution of CVD in India differs from that in developed countries, with a substantial proportion of events occurring in younger individuals (35-64 years), resulting in greater years of life lost and economic impact [25]. This premature CVD burden reflects both genetic susceptibilities and early exposure to risk factors. Indians appear to develop atherosclerosis at younger ages and with less severe traditional risk factor burden than Western populations, suggesting additional environmental or genetic contributions.

Socioeconomic disparities in CVD burden are evident in India, though the pattern differs from that in developed countries. While CVD was historically considered a disease of affluence in India, recent evidence suggests an increasing burden among lower socioeconomic groups, who face greater exposure to risk factors, including air pollution, tobacco use, and psychosocial stress, combined with limited access to preventive and therapeutic healthcare [25].

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

The healthcare system in India faces significant challenges in addressing the CVD epidemic. Limited availability of specialised cardiac care, particularly in rural areas, high out-of-pocket healthcare costs, and inadequate health insurance coverage result in delayed diagnosis, suboptimal treatment, and poor outcomes for many CVD patients. These limitations in the healthcare system also constrain the capacity to identify and protect populations vulnerable to cardiovascular effects induced by air pollution.

Understanding the epidemiology of CVD in India is essential for contextualising the potential impact of air pollution on cardiovascular health. The high and increasing prevalence of CVD, combined with severe air pollution exposure, suggests that ambient particulate matter may contribute substantially to India's cardiovascular disease burden. However, quantifying this contribution requires rigorous epidemiological research that accounts for the complex interplay of traditional risk factors, air pollution exposure, and population-specific susceptibilities.

4. Evidence Linking Air Pollution to Cardiovascular Disease in India

4.1 Epidemiological Studies from Delhi

Delhi has been the focus of the limited epidemiological research on air pollution and health in India, reflecting both the severity of pollution in the capital and the availability of health and environmental data. Two pioneering studies in the late 1990s provided initial evidence of associations between air pollution and health in Delhi, though cardiovascular endpoints were not the primary focus [2,4].

Alberini and colleagues conducted a comprehensive study examining the health effects of air pollution in Delhi using data from the mid-1990s [2]. While the study primarily focused on respiratory symptoms and illnesses. It provided important evidence that Delhi residents experienced substantial health impacts from air pollution exposure.

A companion study by Cropper and colleagues specifically examined the health benefits of air pollution control in Delhi [4]. Using time-series analysis, the researchers investigated associations between daily variations in air pollution levels and mortality. The study found significant associations between particulate matter and total mortality, with estimated effects comparable to or larger than those reported in studies from developed countries. While the analysis did not separately examine cardiovascular mortality, the findings suggested that air pollution contributed to the overall mortality burden in Delhi.

More recently, a pilot study in Mandi-Gobindgarh, Punjab, specifically examined air pollution and cardiovascular health [1]. This industrial town experiences severe air pollution from steel rolling mills and other industries. The study assessed cardiovascular risk factors and health status in relation to air pollution exposure and found elevated blood pressure and other cardiovascular

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

abnormalities in exposed populations compared with control subjects. While limited in scope, this study provided important preliminary evidence of the effects of air pollution on CVD in an Indian population.

The Delhi studies, while groundbreaking, have important limitations. Most did not specifically examine cardiovascular endpoints, limiting direct evidence on air pollution-CVD associations. The studies relied on limited monitoring data and could not assess exposure at the individual level. Furthermore, the studies did not adequately control for potential confounding factors, including traditional cardiovascular risk factors, socioeconomic status, and indoor air pollution.

4.2 Evidence from Mumbai and Other Cities

Evidence from cities other than Delhi remains extremely limited. A case study in Mumbai examined PM10 levels and their health impacts, documenting elevated particulate levels and associated respiratory effects [3]. The study highlighted the public health burden of air pollution in Mumbai but did not specifically investigate cardiovascular outcomes.

Studies in other Indian cities have primarily focused on characterising air quality rather than examining health effects [5]. While these studies document severe particulate pollution in cities including Kolkata, Chennai, Bangalore, and Hyderabad, epidemiological investigations of cardiovascular health impacts are largely absent from the published literature.

The absence of cardiovascular-specific epidemiological studies in most Indian cities represents a critical evidence gap. Given the substantial regional variations in pollution sources, levels, and composition, as well as differences in population characteristics and healthcare access, findings from Delhi cannot be directly extrapolated to other regions. City-specific studies are needed to characterise air pollution-CVD associations across India's diverse urban environments.

4.3 Limitations of Current Evidence

The evidence base linking air pollution to cardiovascular disease in India faces several critical limitations. First, the number of studies is extremely small, with most focusing on respiratory rather than cardiovascular outcomes. Second, available studies have relied on limited air quality monitoring data. Third, most studies have used ecological or time-series designs that cannot establish individual-level exposure-response relationships or adequately control for confounding factors.

Fourth, studies have not examined specific cardiovascular endpoints such as myocardial infarction, stroke, heart failure, or arrhythmias, limiting understanding of which cardiovascular conditions are most affected by air pollution in Indian populations. Fifth, studies have not investigated effect modification by traditional cardiovascular risk factors, genetic susceptibilities, or socioeconomic characteristics, precluding identification of vulnerable subgroups.

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Sixth, the lack of long-term cohort studies prevents assessment of chronic exposure effects on cardiovascular disease incidence and progression. Seventh, studies have not examined the relative contributions of different particulate matter size fractions (PM2.5 vs. PM10) or chemical components to cardiovascular effects. Finally, studies have not assessed the potential confounding or modifying effects of indoor air pollution exposure, which is substantial in India due to widespread biomass fuel use [22,29].

These limitations underscore the urgent need for rigorous, well-designed epidemiological studies to characterise air pollution-CVD associations in Indian populations. Such studies should employ individual-level exposure assessment, examine specific cardiovascular endpoints, investigate vulnerable subgroups, and account for the complex exposure environment in India.

5. Biological Pathways Linking Air Pollution to Cardiovascular Disease

While epidemiological evidence from India remains limited, extensive mechanistic research from other countries has elucidated biological pathways through which particulate air pollution induces cardiovascular injury. Understanding these mechanisms is essential for interpreting epidemiological findings, identifying vulnerable populations, and developing targeted interventions. The primary pathways related to CVD and air pollution are oxidative stress, systemic inflammation, autonomic nervous system dysfunction, and enhanced coagulation and thrombosis [12,13,20,43].

5.1 Oxidative Stress

Oxidative stress is primarily associated with particulate matter exposure and cardiovascular injury [31,32,45]. Inhaled particles can generate reactive oxygen species (ROS) through multiple pathways, including direct chemical redox activity of particle constituents (particularly transition metals and organic compounds), activation of cellular oxidant-generating systems (NADPH oxidase, mitochondrial electron transport), and depletion of antioxidant defences [31,32].

Experimental studies have indicated that particulate matter exposure induces rapid increases in ROS generation in the lungs and cardiovascular system [53]. These reactive species can directly damage cellular macromolecules, including lipids, proteins, and DNA, leading to cellular dysfunction and death [31]. Oxidative modification of low-density lipoprotein (LDL) promotes atherosclerosis development and progression [41,50]. Oxidative stress also activates redox-sensitive transcription factors, including nuclear factor-kappa B (NF-kB), which regulates expression of pro-inflammatory genes [55].

The cardiovascular system is particularly vulnerable to oxidative injury due to high metabolic activity, abundant lipid content, and limited regenerative capacity [45]. Oxidative stress impairs endothelial function by reducing nitric oxide bioavailability. The nitrous oxide is a critical

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

regulator of vascular tone, platelet function, and inflammatory cell adhesion [12,45]. Endothelial dysfunction represents an early step in atherosclerosis development and contributes to acute cardiovascular events [30].

Mitochondrial dysfunction induced by particulate matter exposure represents another important implication of oxidative stress [42]. Particles and their constituents can impair mitochondrial electron transport, reduce ATP production, and trigger apoptotic pathways, compromising cardiac and vascular cell function [42]. These effects may be particularly relevant for individuals with pre-existing cardiovascular disease, whose cardiac mitochondria may have limited reserve capacity.

Individual susceptibility to oxidative stress varies based on genetic polymorphisms in antioxidant enzymes, dietary antioxidant intake, and pre-existing conditions that increase oxidative burden [48,54]. Diabetic individuals, for example, experience elevated baseline oxidative stress and may be particularly vulnerable to additional oxidative injury from air pollution exposure [39,47].

5.2 Systemic Inflammation

Systemic inflammation represents a key pathway linking particulate air pollution to cardiovascular disease [21,28,49]. Inhalation of particles triggers inflammatory responses in the lungs, characterised by the activation of alveolar macrophages, the recruitment of inflammatory cells, and the release of pro-inflammatory mediators [28,34,35]. These mediators, including interleukin-6 (IL-6), tumour necrosis factor-alpha (TNF-alpha), and C-reactive protein (CRP), enter the systemic circulation and promote cardiovascular injury [21,46,49].

Epidemiological studies have documented associations between particulate air pollution exposure and elevated circulating levels of inflammatory biomarkers [21,38,49]. Experimental studies have confirmed that particulate exposure induces systemic inflammation, with effects detectable within hours to days of exposure [37,38]. The magnitude of inflammatory response varies with particle dose, composition, and individual susceptibility factors [21,40].

Systemic inflammation contributes to cardiovascular disease through multiple mechanisms. Inflammatory mediators promote endothelial activation and dysfunction, thereby increasing the expression of adhesion molecules that facilitate the recruitment of inflammatory cells to the vascular wall [51]. Inflammation accelerates atherosclerosis progression by promoting lipid accumulation, smooth muscle cell proliferation, and plaque instability [9,50]. Inflammatory cytokines also induce hepatic production of acute-phase proteins, including fibrinogen and CRP, which contribute to thrombotic risk [44].

The inflammatory response to particulate matter may be particularly pronounced in individuals with pre-existing inflammatory conditions, including diabetes, obesity, and established cardiovascular disease [47]. These populations may experience heightened inflammatory responses to air pollution, increasing their vulnerability to cardiovascular events.

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

The source of systemic inflammation following particulate inhalation remains debated [20]. Inflammation may originate primarily from pulmonary responses, with inflammatory mediators spilling over into systemic circulation. Alternatively, ultrafine particles may translocate across the alveolar-capillary barrier and directly interact with cardiovascular tissues, triggering local inflammatory responses [10,11]. Both mechanisms may contribute, with their relative importance potentially varying with particle size and composition.

5.3 Autonomic Nervous System Dysfunction

Autonomic nervous system dysfunction represents another important mechanism which linking air pollution to cardiovascular effects [13,18,37]. Particulate matter exposure has been associated with altered heart rate variability (HRV), a marker of autonomic balance that predicts cardiovascular risk [18,37]. Reduced HRV, indicating decreased parasympathetic or increased sympathetic activity, has been observed in association with elevated particulate pollution in multiple studies [18,37].

The mechanisms underlying the autonomic effects of air pollution remain incompletely understood [20]. Particle deposition in the lungs may stimulate pulmonary sensory receptors, triggering reflex changes in autonomic outflow [35]. Alternatively, systemic inflammation and oxidative stress may directly affect autonomic control centres in the central nervous system or peripheral autonomic nerves [12].

Autonomic dysfunction can contribute to cardiovascular events through multiple pathways. Increased sympathetic activity raises blood pressure, heart rate, and myocardial oxygen demand while reducing coronary blood flow [13]. Reduced parasympathetic activity diminishes protective vagal influences on the heart, increasing susceptibility to arrhythmias [13]. These effects may be particularly hazardous for individuals with underlying cardiac disease and limited coronary passage. Studies have documented associations between particulate air pollution and cardiac arrhythmias, including ventricular ectopy and atrial fibrillation [24]. These associations may reflect autonomic imbalance, though direct effects of inflammation and oxidative stress on cardiac electrophysiology may also contribute [17]. Elevated blood pressure is another potential consequence of autonomic dysfunction induced by air pollution [16,33]. While evidence for acute blood pressure effects of particulate exposure has been inconsistent, some studies have documented positive associations, particularly in susceptible populations [16]. Chronic exposure to air pollution may contribute to the development of hypertension through sustained autonomic imbalance, vascular remodelling, and endothelial dysfunction [33].

5.4 Coagulation and Thrombosis

Enhanced coagulation and thrombosis represent critical mechanisms linking air pollution to acute cardiovascular events, particularly myocardial infarction and stroke [30,44]. Particulate matter

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

exposure has been associated with increased plasma fibrinogen, enhanced platelet activation, accelerated coagulation, and impaired fibrinolysis [30,38,44].

Experimental studies have demonstrated that particulate exposure induces a prothrombotic state characterised by increased circulating coagulation factors, shortened clotting times, and enhanced thrombus formation [44]. These effects appear to be mediated by pulmonary inflammation, with IL-6 playing a central role in upregulating hepatic synthesis of procoagulant proteins [44]. Direct effects of particles on platelet function and endothelial tissue factor expression may also contribute [38].

The prothrombotic effects of air pollution are particularly relevant for individuals with vulnerable atherosclerotic plaques. Inflammation and oxidative stress induced by particulate exposure may destabilise plaques, increasing the risk of rupture [30]. When plaque rupture occurs in the setting of enhanced coagulation, the likelihood of occlusive thrombus formation and acute cardiovascular events increases substantially [30].

Epidemiological studies have documented associations between particulate air pollution and acute myocardial infarction, with effects apparent within hours to days of exposure [30]. These temporal patterns are consistent with thrombotic mechanisms, which can rapidly precipitate acute events in susceptible individuals. The magnitude of thrombotic risk associated with air pollution appears comparable to that of other recognized triggers of acute cardiovascular events, including physical exertion and emotional stress [30].

Individual susceptibility to the prothrombotic effects of air pollution varies according to baseline coagulation status, anticoagulant medication use, and the presence of vulnerable atherosclerotic plaques [30]. Patients with established coronary artery disease, prior myocardial infarction, or multiple cardiovascular risk factors may be at particularly high risk for pollution-triggered thrombotic events [38].

6. Regional Disparities in Air Pollution and Cardiovascular Risk

India's vast geographic, climatic, and socioeconomic diversity results in substantial regional disparities in air pollution levels, sources, and composition. These disparities have important implications for cardiovascular health impacts and necessitate region-specific approaches to exposure assessment, epidemiological research, and pollution control.

6.1 Delhi and North India

Delhi and the broader North Indian region experience the most severe air pollution in India, with particulate levels among the highest globally [2,4,5]. Multiple factors contribute to this extreme pollution burden. The topography and meteorology of the Indo-Gangetic Plain limit pollutant dispersion, particularly during winter months when temperature inversions trap pollutants near the surface [5]. The region's dense population, heavy vehicular traffic, and industrial activity generate

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

high levels of emissions. Agricultural burning of crop residues, particularly rice and wheat stubble, contributes to severe episodic pollution events [5].

Delhi's air pollution is characterised by high concentrations of both PM10 and PM2.5, with significant contributions from vehicular emissions, industrial sources, construction dust, and biomass burning [2,4,5]. The city's vehicle fleet has expanded rapidly, and despite some pollution control measures, such as CNG conversion of public transport, vehicular emissions remain a dominant source [4]. Industrial areas within and surrounding Delhi contribute additional pollution, particularly from small-scale industries with limited emission controls [5].

The cardiovascular health implications of Delhi's severe air pollution are likely substantial, though direct evidence remains limited [1,2,4]. The high pollution levels, combined with increasing CVD prevalence and traditional risk factors, suggest that air pollution may contribute significantly to cardiovascular morbidity and mortality in the region. The limited epidemiological evidence from Delhi indicates associations between air pollution and mortality, though cardiovascular-specific effects require further investigation [2,4].

Other cities in North India, including Kanpur, Lucknow, and Agra, also experience severe particulate pollution, though monitoring data and health studies are limited [5]. Festival-related pollution events, such as those documented during Diwali in Lucknow, create acute exposure peaks with potential cardiovascular implications [26]. Rural areas in North India face substantial indoor air pollution from biomass fuel use, which may interact with pollution to increase cardiovascular risk [22,29].

6.2 Kolkata and East India

Kolkata, the capital of West Bengal and a major metropolitan centre in eastern India, faces substantial air pollution from vehicular traffic, industrial emissions, and domestic sources [5]. The city's dense population, narrow streets, and limited green spaces exacerbate exposure to pollution. Monitoring studies have documented high PM10 and TSP levels, with significant seasonal variation and spatial heterogeneity [5].

Particulate pollution in Kolkata shows distinct seasonal patterns, with the highest levels during winter months when meteorological conditions limit dispersion and domestic biomass burning increases [5]. The post-monsoon period also experiences elevated pollution due to festival-related activities. Industrial areas of Kolkata experience higher pollution levels than residential areas, though all areas exceed air quality standards [5].

The composition of particulate matter in Kolkata reflects contributions from vehicular emissions, industrial sources, and biomass combustion [5]. The city's industrial base includes jute mills, chemical industries, and small-scale manufacturing, all of which contribute to air pollution. The cardiovascular health impacts of air pollution in Kolkata have not been systematically studied,

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

which represents a critical knowledge gap. The city's high pollution levels, combined with increasing CVD prevalence, suggest substantial potential health impacts.

6.3 Mumbai and West India

Mumbai, India's financial capital and most populous city, experiences severe air pollution, though levels are generally lower than in Delhi and Kolkata [3,5]. The city's coastal location and sea breezes help disperse pollutants, but high population density, intensive vehicular traffic, and industrial activity result in substantial exposure to pollutants [3].

A case study in Mumbai documented PM10 levels frequently exceeding air quality standards, with significant health impacts [3]. Monitoring data from Mumbai shows spatial variation in pollution levels, with higher concentrations in industrial areas and along major traffic corridors [5]. Vehicular emissions are the primary source of air pollution in Mumbai, with the city's large vehicle fleet and traffic congestion contributing to elevated levels of particulate matter and gaseous pollutants [5].

The cardiovascular health implications of Mumbai's air pollution remain poorly characterised. The city's large population, rising CVD prevalence, and substantial exposure to air pollution suggest that air pollution may contribute significantly to the cardiovascular disease burden. Other cities in western India, including Pune and Ahmedabad, also experience significant air pollution, though monitoring data and health studies are limited [5]. The region's industrial activities, including automobile manufacturing, chemical production, and textile industries, contribute to local and regional air pollution.

6.4 South India

Cities in southern India generally experience lower air pollution levels than those in the north, though levels still frequently exceed air quality standards [5]. Bangalore, Chennai, and Hyderabad, the region's major metropolitan centres, face increasing pollution from vehicular traffic and industrial growth, though meteorological conditions and lower population densities result in better dispersion of pollutants than in northern cities [5].

Bangalore's air pollution is primarily driven by vehicular emissions, with the city's rapidly expanding vehicle fleet and traffic congestion contributing to elevated levels of particulate and gaseous pollutants [5]. The city's substantial population growth imposes increasing pressure on transportation infrastructure and air quality. Chennai faces pollution from vehicular traffic, industrial sources including petrochemical facilities and power plants, and port activities [5].

Hyderabad experiences air pollution from vehicular emissions, industrial activities, and construction [5]. The cardiovascular health impacts of air pollution in southern Indian cities have not been systematically studied. Lower pollution levels than in northern cities may result in smaller absolute health impacts, though effects may still be substantial given the large populations

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

exposed. The region's increasing CVD prevalence and changing risk factor profiles necessitate research on the relationships between air pollution and cardiovascular health.

6.5 Rural versus Urban Disparities

Substantial disparities exist between rural and urban areas in both air pollution exposure and cardiovascular disease burden [5,25]. Urban areas experience higher air pollution levels due to vehicular traffic, industrial emissions, and higher population densities [5]. Rural areas, while experiencing lower pollution, face substantial indoor air pollution from biomass fuel combustion for cooking and heating [22,29].

Indoor air pollution from biomass fuels represents a major health hazard in rural India, with women and children experiencing particularly high exposures [22,29]. While indoor air pollution has primarily been studied in relation to respiratory diseases, emerging evidence suggests cardiovascular effects [22]. The interaction between indoor and air pollution exposures in semi-urban areas, where both sources contribute substantially, remains poorly understood.

Cardiovascular disease prevalence shows an urban-rural gradient, with higher rates in cities [25]. However, rural CVD burden is increasing, and the contribution of air pollution (both indoor and ambient) to this trend requires investigation. The rural-urban disparity in air pollution and CVD has important policy implications. Urban areas require interventions to reduce air pollution from vehicular and industrial sources, while rural areas need programs to promote clean cooking fuels and technologies to reduce exposure to indoor air pollution.

7. Vulnerable Populations

Certain population subgroups face heightened vulnerability to cardiovascular effects of air pollution due to increased exposure, enhanced biological susceptibility, or limited capacity to avoid or mitigate exposure [13,27,43]. Identifying and protecting these vulnerable populations is essential for public health priorities in India.

Elderly individuals are more vulnerable to cardiovascular effects of air pollution due to age-related changes in cardiovascular function, a higher prevalence of pre-existing cardiovascular disease, and potentially diminished physiological reserve [27]. Studies from other countries have consistently documented stronger associations between air pollution and mortality in elderly populations [9,27]. The elderly population in India is growing rapidly, increasing the number of individuals at heightened risk.

Individuals with pre-existing cardiovascular disease represent a particularly vulnerable population [27,52]. Patients with coronary artery disease, heart failure, prior myocardial infarction, or arrhythmias have limited cardiac reserve and may be unable to bear the additional stress imposed by air pollution exposure [27,52]. Studies have documented increased risk of hospital readmissions and mortality among cardiac patients exposed to elevated air pollution [27,52].

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Diabetic individuals may be more susceptible to the cardiovascular effects of air pollution [39,47]. Diabetes is associated with endothelial dysfunction, increased oxidative stress, systemic inflammation, and prothrombotic state—all mechanisms through which air pollution exerts cardiovascular effects [39,47]. Studies have documented stronger associations between air pollution and cardiovascular outcomes in diabetic compared to non-diabetic individuals [39,47]. Given India's large and growing diabetic population, this represents a substantially vulnerable group.

Individuals of lower socioeconomic status face multiple vulnerabilities to the cardiovascular effects of air pollution [19]. Lower-income populations often experience greater exposure to pollution due to residence in more polluted areas, proximity to traffic and industrial sources, and occupational exposures [19,23]. They also have a higher prevalence of cardiovascular risk factors, limited access to healthcare, and reduced capacity to avoid or mitigate exposure to pollution [19]. Traffic police and other outdoor workers face particularly high exposures [23].

Women may face specific vulnerabilities related to indoor air pollution exposure from cooking with biomass fuels [22,29]. While indoor air pollution has been primarily studied in relation to respiratory diseases, potential cardiovascular effects warrant investigation. Women also face barriers to healthcare access in many parts of India, potentially limiting their ability to receive treatment for pollution-related cardiovascular effects.

Individuals with genetic susceptibilities may be more vulnerable to cardiovascular effects of air pollution [48]. Genetic polymorphisms affecting antioxidant enzymes, inflammatory mediators, and coagulation factors may modify individual responses to particulate exposure [48]. However, research on gene-environment interactions in air pollution health effects remains limited, particularly in Indian populations.

Populations with multiple risk factors face compounded vulnerability. Individuals with combinations of advanced age, pre-existing cardiovascular disease, diabetes, low socioeconomic status, and high pollution exposure face a particularly high risk of pollution-induced cardiovascular effects. Identifying and protecting these multiply vulnerable populations requires integrated approaches addressing both pollution exposure and underlying health conditions.

8. Policy Implications and Gaps

The emerging evidence on cardiovascular vulnerability to air pollution in India has important implications for public health policy, though substantial gaps in knowledge and implementation capacity constrain effective action.

Air quality monitoring requires urgent strengthening. The current monitoring network is inadequate in terms of spatial coverage, temporal continuity, and the measurement of health-relevant pollutants, particularly PM2.5 [5]. Expanding the monitoring network to cover more cities, establishing monitors in diverse microenvironments (residential, traffic, industrial), ensuring continuous operation and quality control, and implementing PM2.5 monitoring are

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

essential priorities. Enhanced monitoring will support epidemiological research, inform the public about air quality, and enable evaluation of pollution control interventions.

Epidemiological research on air pollution and cardiovascular health in India requires substantial expansion. Priority research needs include: (1) time-series and case-crossover studies examining associations between short-term pollution fluctuations and acute cardiovascular events (myocardial infarction, stroke, heart failure exacerbations, arrhythmias, cardiovascular mortality); (2) cohort studies investigating long-term exposure effects on cardiovascular disease incidence and progression; (3) panel studies in vulnerable populations (elderly, cardiac patients, diabetics) examining subclinical cardiovascular effects; (4) studies investigating effect modification by traditional risk factors and genetic susceptibilities; and (5) health impact assessments quantifying the cardiovascular disease burden attributable to air pollution in India.

Pollution source control represents the most effective approach to protecting cardiovascular health. Priority interventions include: (1) strengthening and enforcing vehicular emission standards; (2) improving fuel quality; (3) expanding public transportation and promoting non-motorised transport; (4) implementing industrial emission controls; (5) controlling construction dust; (6) managing agricultural burning; and (7) promoting clean cooking fuels and technologies to reduce indoor air pollution [4,5,19]. The effectiveness of these interventions requires rigorous evaluation.

Healthcare system preparedness for cardiovascular effects related to air pollution requires attention. Healthcare providers need education about air pollution as a cardiovascular risk factor and guidance on advising vulnerable patients during high-pollution episodes. Health systems need the capacity to monitor and respond to increases in cardiovascular events driven by pollution. Integration of air quality information into clinical decision-making and public health surveillance systems could enhance protection of vulnerable populations.

Public awareness and communication about air pollution and cardiovascular health remain limited in India. Public education campaigns highlighting cardiovascular risks of air pollution, particularly for vulnerable populations, could promote support for pollution control policies. Real-time air quality information systems accessible across multiple platforms could enable individuals to reduce exposure during high-pollution episodes.

Critical gaps that constrain effective action include: (1) limited air quality monitoring data; (2) paucity of epidemiological evidence on air pollution-CVD associations in Indian populations; (3) lack of exposure assessment tools appropriate for Indian contexts; (4) limited understanding of population-specific susceptibilities; (5) inadequate evaluation of pollution control intervention effectiveness; (6) weak enforcement of existing air quality standards; (7) limited public awareness of air pollution health risks; and (8) insufficient healthcare system capacity to address pollution-related health effects.

9. Conclusions

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

Air pollution represents an emerging and underappreciated cardiovascular risk factor in India. The convergence of severe particulate pollution and rapidly increasing cardiovascular disease burden creates a critical public health challenge. While direct epidemiological evidence from India remains limited, the extreme pollution levels, established biological mechanisms linking particulate matter to cardiovascular injury, and consistent findings from international studies strongly suggest that air pollution contributes substantially to cardiovascular morbidity and mortality in India.

Urban India experiences some of the highest particulate air pollution levels globally, with PM10 concentrations frequently exceeding 200 μg/m3 in major cities. These extreme levels reflect contributions from vehicular emissions, industrial activities, biomass combustion, and other sources, compounded by meteorological conditions that limit pollutant dispersion. Substantial regional disparities exist, with northern cities experiencing the most severe pollution, though all major urban cities face significant air quality challenges.

The biological pathways linking particulate matter to cardiovascular disease are well-established and include oxidative stress, systemic inflammation, autonomic nervous system dysfunction, and enhanced coagulation and thrombosis. These mechanisms can trigger acute cardiovascular events in susceptible individuals and contribute to chronic disease development and progression. The consistency of these pathways across populations suggests that Indian populations exposed to high particulate levels face substantial cardiovascular risk.

Vulnerable populations, including the elderly, individuals with pre-existing cardiovascular disease, diabetics, and those of lower socioeconomic status, face a heightened risk of pollution-induced cardiovascular effects. Protecting these populations requires targeted interventions addressing both pollution exposure and underlying health conditions. The large and growing size of vulnerable populations in India amplifies the public health significance of air pollution-cardiovascular health relationships.

Future research priorities include: (1) establishing long-term cohort studies to investigate chronic exposure effects; (2) conducting time-series and case-crossover studies of acute cardiovascular events; (3) investigating population-specific susceptibilities and effect modifiers; (4) characterizing exposure patterns using personal monitoring and modeling approaches; (5) examining the relative toxicity of different particulate matter components; (6) assessing interactions between ambient and indoor air pollution; (7) evaluating the effectiveness of pollution control interventions; and (8) conducting health impact assessments to quantify the cardiovascular disease burden attributable to air pollution.

In conclusion, air pollution represents a significant and growing threat to cardiovascular health in India. The extreme pollution levels, established biological mechanisms, and emerging epidemiological evidence indicate that air pollution contributes substantially to India's cardiovascular disease burden. Protecting cardiovascular health requires urgent action to reduce exposure to pollution through effective source controls, enhance scientific understanding through

rigorous research, and strengthen health system capacity to identify and protect vulnerable populations.

References

1. Nautiyal J, Garg ML, Kumar M, et al. Air Pollution and Cardiovascular Health in Mandi-Gobindgarh, Punjab, India—A Pilot Study. Int J Environ Res Public Health. 2007;4(4):302-311. doi:10.3390/ijerph200704040002

2. Alberini A, Cropper ML, Simon NB, et al. The Health Effects of Air Pollution in Delhi, India. World Bank Policy Research Working Paper 1860. Washington, DC: World Bank; 1997. doi:10.1596/1813-9450-1860

3. Joseph AE, Sawant AD, Srivastava A. PM(10) and its impacts on health—a case study in Mumbai. Int J Environ Health Res. 2003;13(2):207-214. doi:10.1080/0960312031000098107

4. Cropper ML, Simon NB, Alberini A, et al. The Health Benefits of Air Pollution Control in Delhi. Am J Agric Econ. 1997;79(5):1625-1629. doi:10.2307/1244393

5. Kandlikar M, Ramachandran G. The Causes and Consequences of Particulate Air Pollution in Urban India: A Synthesis of the Science. Annu Rev Energy Environ. 2000;25:629-684. doi:10.1146/annurev.energy.25.1.629

6. Dockery DW. Epidemiologic Evidence of Cardiovascular Effects of Particulate Air Pollution. Environ Health Perspect. 2001;109 Suppl 4:483-486. doi:10.2307/3454657

7. Kunzli N, Tager IB. Air pollution: from lung to heart. Swiss Med Wkly. 2005;135(47-48):697-702. doi:10.4414/smw.2005.11025

8. Mohanraj R, Azeez PA. Health effects of airborne particulate matter and the Indian scenario. Curr Sci. 2004;87(6):741-748.

9. Pope CA III, Burnett RT, Thurston GD, et al. Cardiovascular Mortality and Long-Term Exposure to Particulate Air Pollution: Epidemiological Evidence of General Pathophysiological Pathways of Disease. Circulation. 2004;109(1):71-77. doi:10.1161/01.CIR.0000108927.80044.7F

10. Delfino RJ, Sioutas C, Malik S. Potential role of ultrafine particles in associations between airborne particle mass and cardiovascular health. Environ Health Perspect. 2005;113(8):934-946. doi:10.1289/ehp.7938

11. Schulz H, Harder V, Ibald-Mulli A, et al. Cardiovascular effects of fine and ultrafine particles. J Aerosol Med. 2005;18(1):1-22. doi:10.1089/jam.2005.18.1

12. Bhatnagar A. Environmental Cardiology: Studying Mechanistic Links Between Pollution and Heart Disease. Circ Res. 2006;99(7):692-705. doi:10.1161/01.RES.0000243586.99701.CF

13. Brook RD, Franklin BA, Cascio WE, et al. Air Pollution and Cardiovascular Disease: A Statement for Healthcare Professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. Circulation. 2004;109(21):2655-2671. doi:10.1161/01.CIR.0000128587.30041.C8

14. Brook RD. Is air pollution a cause of cardiovascular disease? Updated review and controversies. Rev Environ Health. 2007;22(2):115-137. doi:10.1515/REVEH.2007.22.2.115

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

15. Dockery DW, Stone PH. Cardiovascular Risks from Fine Particulate Air Pollution. N Engl J Med. 2007;356(5):511-513. doi:10.1056/NEJMe068274

16. Auchincloss AH, Diez Roux AV, Dvonch JT, et al. Associations between Recent Exposure to Ambient Fine Particulate Matter and Blood Pressure in the Multi-Ethnic Study of Atherosclerosis (MESA). Environ Health Perspect. 2008;116(4):486-491. doi:10.1289/ehp. 10899

17. Zareba W, Nomura A, Couderc JP. Cardiovascular Effects of Air Pollution: What to Measure in ECG? Environ Health Perspect. 2001;109 Suppl 4:533-538. doi:10.2307/3454665

18. Chan CC, Chuang KJ, Su TC, et al. Association between nitrogen dioxide and heart rate variability in a susceptible population. Eur J Prev Cardiol. 2005;12(6):580-586. doi:10.1097/01.hjr.0000185975.44265.2e

19. Kjellstrom T, Lodh M, McMichael T, et al. Air and Water Pollution: Burden and Strategies for Control. In: Jamison DT, Breman JG, Measham AR, et al., eds. Disease Control Priorities in Developing Countries. 2nd ed. Washington, DC: World Bank; 2006. chap. 43.

20. Kaufman JD. Air pollution and mortality: are we closer to understanding the how? Am J Respir Crit Care Med. 2007;176(4):325-326. doi:10.1164/rccm.200705-647ED

21. Zeka A, Sullivan JR, Vokonas PS, et al. Inflammatory markers and particulate air pollution: characterizing the pathway to disease. Int J Epidemiol. 2006;35(5):1347-1354. doi:10.1093/ije/dyl132

22. Smith KR. Indoor air pollution in developing countries: recommendations for research. Indoor Air. 2002;12(3):198-207. doi:10.1034/j.1600-0668.2002.01137.x

23. Ingle ST, Pachpande BG, Wagh ND, et al. Exposure to Vehicular Pollution and Respiratory Impairment of Traffic Policemen in Jalgaon City, India. Ind Health. 2005;43(4):656-662. doi:10.2486/indhealth.43.656

24. Ljungman P, Berglind N, Holmgren C, et al. Rapid effects of air pollution on ventricular arrhythmias. Eur Heart J. 2008;29(23):2894-2901. doi:10.1093/eurheartj/ehn463

25. Gupta R, Joshi PP, Mohan V, et al. Epidemiology and causation of coronary heart disease and stroke in India. Heart. 2008;94(1):16-26. doi:10.1136/hrt.2007.132951

26. Barman SC, Singh R, Negi MPS, et al. Ambient air quality of Lucknow City (India) during use of fireworks on Diwali Festival. Environ Monit Assess. 2008;137(1-3):495-504. doi:10.1007/s10661-007-9784-1

27. Medina-Ramon M, Goldberg RJ, Melly SJ, et al. Residential exposure to traffic-related air pollution and survival after heart failure. Environ Health Perspect. 2008;116(4):481-485. doi:10.1289/ehp.10918

28. van Eeden SF, Tan WC, Suwa T, et al. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants (PM10). Am J Respir Crit Care Med. 2001;164(5):826-830. doi:10.1164/ajrccm.164.5.2010160

29. Bruce N, Perez-Padilla R, Albalak R. Indoor air pollution in developing countries: a major environmental and public health challenge. Bull World Health Organ. 2000;78(9):1078-1092. doi:10.1590/S0042-96862000000900004

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

30. Mills NL, Tornqvist H, Robinson SD, et al. Air Pollution and Atherothrombosis. Inhal Toxicol. 2007;19(Suppl 1):81-89. doi:10.1080/08958370701495170

31. Risom L, Moller P, Loft S. Oxidative stress-induced DNA damage by particulate air pollution. Mutat Res. 2005;592(1-2):119-137. doi:10.1016/j.mrfmmm.2005.06.012

32. Valavanidis A, Fiotakis K, Vlachogianni T. Airborne particulate matter and human health: toxicological assessment and importance of size and composition of particles for oxidative damage and carcinogenic mechanisms. J Environ Sci Health C. 2008;26(4):339-362. doi:10.1080/10590500802494538

33. Brook RD. Why Physicians Who Treat Hypertension Should Know More About Air Pollution. J Clin Hypertens. 2007;9(8):629-635. doi:10.1111/j.1524-6175.2007.07187.x

34. MacNee W, Li XY, Gilmour PS, et al. Systemic Effect of Particulate Air Pollution. Inhal Toxicol. 2000;12(Suppl 3):233-244. doi:10.1080/08958378.2000.11463218

35. Donaldson K, Stone V, Seaton A, et al. Ambient particle inhalation and the cardiovascular system: potential mechanisms. Environ Health Perspect. 2001;109(Suppl 4):523-527. doi:10.1289/ehp.01109s4523

36. Dockery DW, Stone PH. Cardiovascular Risks from Fine Particulate Air Pollution. N Engl J Med. 2007;356(5):511-513. doi:10.1056/NEJMe068274

37. Chuang KJ, Chan CC, Su TC, et al. The Effect of Urban Air Pollution on Inflammation, Oxidative Stress, Coagulation, and Autonomic Dysfunction in Young Adults. Am J Respir Crit Care Med. 2007;176(4):370-376. doi:10.1164/rccm.200611-1627OC

38. Delfino RJ, Staimer N, Tjoa T, et al. Circulating biomarkers of inflammation, antioxidant activity, and platelet activation are associated with primary combustion aerosols in subjects with coronary artery disease. Environ Health Perspect. 2008;116(7):898-906. doi:10.1289/ehp.11189

39. Liu L, Ruddy TD, Dalipaj MM, et al. Influence of personal exposure to particulate air pollution on cardiovascular physiology and biomarkers of inflammation and oxidative stress in subjects with diabetes. J Occup Environ Med. 2007;49(3):258-268. doi:10.1097/jom.0b013e31803220ef

40. Scapellato ML, Lotti M. Short-Term Effects of Particulate Matter: An Inflammatory Mechanism? Crit Rev Toxicol. 2007;37(6):461-487. doi:10.1080/10408440701385622

41. Araujo JA, Barajas B, Kleinman MT, et al. Ambient Particulate Pollutants in the Ultrafine Range Promote Early Atherosclerosis and Systemic Oxidative Stress. Circ Res. 2008;102(5):589-596. doi:10.1161/circresaha.107.164970

42. Xia T, Kovochich M, Nel AE. Impairment of mitochondrial function by particulate matter (PM) and their toxic components: implications for PM-induced cardiovascular and lung disease. Front Biosci. 2007;12:1238-1246. doi:10.2741/2142

43. Brook RD, Franklin BA, Cascio WE, et al. Air Pollution and Cardiovascular Disease: A Statement for Healthcare Professionals from the American Heart Association. Circulation. 2004;109(21):2655-2671. doi:10.1161/01.CIR.0000128587.30041.C8

44. Burton A. Particulate matter: the clot thickens: new clues to PM action. Environ Health Perspect. 2007;115(12):A580. doi:10.1289/ehp.115-a580b

ISSN: 0975-3583,0976-2833 VOL 1, ISSUE 1, 2010

45. Bhatnagar A. Environmental Cardiology: Studying Mechanistic Links Between Pollution and Heart Disease. Circ Res. 2006;99(7):692-705. doi:10.1161/01.RES.0000243586.99701.CF

46. Vogel CFA, Sciullo E, Wong P, et al. Induction of proinflammatory cytokines and C-reactive protein in human macrophage cell line U937 exposed to air pollution particulates. Environ Health Perspect. 2005;113(11):1536-1541. doi:10.1289/ehp.8094

47. O'Neill MS, Veves A, Sarnat JA, et al. Air pollution and inflammation in type 2 diabetes: a mechanism for susceptibility. Occup Environ Med. 2007;64(6):373-379. doi:10.1136/oem.2006.030023

48. Chahine T, Baccarelli AA, Litonjua AA, et al. Particulate air pollution, oxidative stress genes, and heart rate variability in an elderly cohort. Environ Health Perspect. 2007;115(11):1617-1622. doi:10.1289/ehp.10318

49. Zeka A, Sullivan JR, Vokonas PS, et al. Inflammatory markers and particulate air pollution: characterizing the pathway to disease. Int J Epidemiol. 2006;35(5):1347-1354. doi:10.1093/ije/dyl132

50. Sun Q, Wang A, Jin X, et al. Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model. JAMA. 2005;294(23):3003-3010. doi:10.1001/jama.294.23.3003

51. Kristovich RL, Knight DA, Long JF, et al. Macrophage-mediated endothelial inflammatory responses to airborne particulates: impact of particulate physicochemical properties. Chem Res Toxicol. 2004;17(9):1204-1211. doi:10.1021/tx049893p

52. von Klot S, Peters A, Aalto P, et al. Ambient Air Pollution Is Associated With Increased Risk of Hospital Cardiac Readmissions of Myocardial Infarction Survivors in Five European Cities. Circulation. 2005;112(20):3073-3079. doi:10.1161/CIRCULATIONAHA.105.548743

53. Gurgueira SA, Lawrence J, Coull BA, et al. Rapid increases in the steady-state concentration of reactive oxygen species in the lungs and heart after particulate air pollution inhalation. Environ Health Perspect. 2002;110(8):749-755. doi:10.1289/ehp. 02110749

54. Kelly FJ. Dietary antioxidants and environmental stress. Proc Nutr Soc. 2004;63(4):579-585. doi:10.1079/PNS2004388

55. Kennedy TP, Ghio AJ, Reed W, et al. Copper-dependent Inflammation and Nuclear Factor-kappaB Activation by Particulate Air Pollution. Am J Respir Cell Mol Biol. 1998;19(3):366-378. doi:10.1165/ajrcmb.19.3.3042