

Characteristics Of Patients With Left Ventricular Free Wall Rupture Related Death In Acute Myocardial Infarction Setting: A Retrospective Analysis Of Patients From A Single Tertiary Cardiac Centre.

¹HARSSHA BASAPPA,²AMEET M KUMAR,^{3*}SACHIN RAO,⁴SADANANDA .K. S,
⁵SANTHOSH .K, ⁶MANJUNATH . C. N.

¹Sri Jayadeva Institute of cardiovascular Sciences and Research , Mysore Branch Associate Professor, Cardiology ,drharshabasappa@gmail.com

²Sri Jayadeva Institute of cardiovascular Sciences and Research , Mysore Branch Assistant Professor, Pediatric Cardiology, doc.ameet22@gmail.com

^{3*}Sri Jayadeva Institute of cardiovascular Sciences and Research , Mysore Branch Assistant Professor, Cardiology, rao.sachin86@gmail.com

⁴Sri Jayadeva Institute of cardiovascular Sciences and Research , Mysore Branch Professor, Head of Department Cardiology ,sadanand_dm@rediffmail.com

⁵ Sri Jayadeva Institute of cardiovascular Sciences and Research , Mysore Branch ,Associate Professor, Cardiology ,drsanthoshk@gmail.com

⁶Sri Jayadeva Institute of cardiovascular Sciences and Research , Bengaluru Director , SJICSR, Bengaluru

Abstract:

Introduction: Left Ventricular free wall rupture (LVFWR) is a lethal and rare complication of acute myocardial infarction (AMI). LVFWR accounts for 8–17% of mortality in STEMI patients. Characteristics, predisposing factors which are vital in preventing this complication are yet to be studied well.

Objective: To study the characteristics and predisposing factors of patients with LVFWR related death in AMI setting.

Methods: All AMI patients, whose death is documented to be due to primarily LVFWR, were included in the study. Patient's demographic data, risk/precipitating factors, clinical characteristics, in-hospital course, revascularization strategies, and angiographic data were collected from our institutional database.

Results: A total of 69 patients with LVFWR related deaths were identified. Mean age of the study subjects was 65.32 years. 36 patients were females, 33 were males. 40% of subjects were in the age group of 61-70 years. Hypertension is the most common associated risk factor seen in 55% patients. Mean duration of presentation since the onset of symptoms is around 23 hours. Thirty patients (43%) presented in a state of cardiogenic shock. Beta blockers could be initiated in just above 20% of patients. Majority of patients (71%) received thrombolysis. Majority had single vessel disease (61% of those who underwent angiogram). LAD was the most common Culprit vessel seen in 8 out of 13 angiograms (61.5%).

Conclusion: Classical risk factors of LVFWR like advanced age, female gender, delayed presentation, presence of hypertension; first acute MI and cardiogenic shock at the time of presentation were well appreciated. Majority of LVFWR patients had single vessel disease predominantly left anterior descending artery disease. Majority had undergone thrombolysis.

Keywords: Left ventricular free wall rupture, acute myocardial infarction, STEMI, cardiogenic shock

Introduction

Left Ventricular free wall rupture (LVFWR) is a lethal and rare complication of acute myocardial infarction (AMI). Its Incidence is 1.3–2.5% in STEMI patients.¹⁻³ LVFWR accounts for 8–17% of mortality in STEMI patients.⁴ It can occur in one of two types: either acute lethal form or a subacute form where a blood clot in rare circumstances seals the defect and results in the formation of a ventricular pseudoaneurysm. Subacute variant occurs in one third of LVFWR patients.

Approximately 50% of patients with LVFWR are diagnosed within 5 days of the AMI, with 90% diagnosed within 2 weeks. Risk factors for the development of free-wall rupture following AMI include age, female gender, hypertension, first MI, and poor coronary artery collateralization. The advent of primary

percutaneous interventions (PCI), when compared to the pre-thrombolytic or the thrombolytic eras, has considerably reduced the rates of LVFWR.⁵

Urgent transthoracic echocardiography (TTE) is the gold standard for definitive diagnosis of LVFWR. Cardiac Magnetic Resonance (CMR) / Multi-Detector Computed Tomography (MDCT) are feasible options for well characterization of Pseudoaneurysm in subacute cases. Urgent surgical repair is still the rule for treatment of LVFWR using pericardial patch closure or less frequently infarctectomy with patch placement and ventricular wall reconstruction. Recently sutureless techniques are being employed more frequently.

LVFWR carries a poor prognosis. However, due to its rare and unpredictable incidence, there is a large lacuna in our understanding of the disease. Not many studies are available in the literature regarding LVFWR. Characteristics, predisposing factors which are vital in preventing this complication are yet to be studied well, as previous Indian studies have included a very small sample size. This study, with a relatively larger sample size would be helpful in understanding the characteristic of the disease better in Indian setup.

Materials and methods:

This is a retrospective observational study. All acute MI (AMI) patients, whose death is documented to be due to primarily LVFWR, were included in the study. Patient’s demographic data, risk/precipitating factors, clinical characteristics, in-hospital course, revascularization strategies, and angiographic data (wherever applicable), were collected from our institutional database between August 2018 and May 2020. Identification of LVFWR was primarily done by 2D Echocardiographic parameters. As an institution protocol, LVFWR is identified on 2D echocardiography by an experienced board certified cardiologist. Useful findings in Echocardiography involve a pericardial collection with signs of cardiac tamponade, including collapse of the right atrium and ventricle in diastole, a dilated inferior vena cava, and marked respiratory variation in mitral and tricuspid valve inflow. **Statistical analysis:**

Descriptive statistical analysis has been carried out in the present study, expressed as percentage and mean.

Results

From the database of patients presenting with AMI between August 2018 and May 2020, 69 patients with LVFWR related deaths were identified. Table I shows the gender distribution of the subjects. Mean age of the study subjects was 65.32 years. 36 patients were females, 33 were males.

Table I. Gender distribution of Study subjects

Gender distribution	N (%)	Mean age (y)
Males	33 (47.8)	63.3
Females	36 (52.1)	67.3
Overall	69	65.3

Table II: Age distribution of study subjects

Age group in years	N(%)
>80	2 (2.8)
71-80	17 (24.6)
61-70	27 (39.1)
</=60	23 (33.33)

Majority of the study subjects i.e. 40% were in the age group of 61-70 years as shown in Table II. Very elderly i.e. > 80 y constituted less than 3% of patients.

Table III: Distribution of studysubjects based on Risk factors

Associated risk factors	N (%)
Past history of IHD	8 (11.5)
Diabetes mellitus	20 (28.9)
Hypertension	38 (55)
Smoking	32 (46.3)
Dyslipidemia	8 (11.5)

As shown in Table III, Hypertension is the most common associated risk factor seen in 55% patients, followed by smoking which is seen in 46% of patients. Diabetes Mellitus is a less frequently seen risk factor in only 20 out of 69 patients (28.9%). Only eight patients (11%) had past history of IHD.

Table IV. Details of clinical presentation

Clinical presentation	Frequency (N) %
MeanDuration between symptom onsetand presentation to the hospital	23.38 hours
MeanTIMI score	5.83
Type of MI:	
AMI site	
-Anterior	44(63.7%)
-Inferior	20 (28.9%)
-Lateral	2 (2.8%)
-Right	0
-Posterior	9 (13%)
Timing of ruptue:	
Early(< 48 h)	40(57.9%)
Late (>48 h)	29 (42%)
Killip class	
I	25 (36.2%)
II	8 (11.5%)
III	6 (8.6%)
IV	30(43.4%)

Table IV shows the details of clinical presentation. Mean duration of presentation since the onset of symptoms is around 23 hours. Forty four patients presented less than 24 h (63.76%) and 25 patients presented after 24 h duration (36.23%). Thirty patients (43%) presented in a state of cardiogenic shock. Eight patient out of 69 (11.5%) had complete heart block. One patient had 2:1 AV block in ECG.

Table V. Investigations

Parameter	Mean value
Laboratory Parameters :	
Creatinine	1.12
Troponin T	1.36
Echo Parameters :	
EF	36.09
LVIDd	4.85

Table V shows relevant investigation findings. Mean EF at the time of presentation was just over 36%.

Table VI. Treatment received.

Treatment	N(%)
Medication:	
• Beta blockers	15 (21.7)
• ACE inhibitors/ ARBs	37 (53.6)
• Statins	69 (100)
Revascularization	
• Thrombolysis	49 (71)
• PTCA	
-Facilitated	3 (4.3)
-Rescue	5 (7.2)
-Primary	4 (5.7)

Table VI shows the treatment received by the patients. Beta blockers could be initiated in just above 20% of patients. Majority of patients (71%) received thrombolysis with streptokinase. Out of which five patients had failed thrombolysis and underwent rescue PTCA. Other 3 patients underwent elective PTCA.

Table VII: Coronary Angiographic data

Coronary Angiogram Findings	N(%)
CAG performed	13
LM	1 (7.6)
SVD	8 (61.5)
DVD	3 (23)
TVD	2 (28.9)
Culprit vessel PTCA	Total- 12
LAD	8 (66.6)
LCX	3 (25)
RCA	1 (8.3)

Table VII shows the coronary angiogram data. Only 13 out of 69 underwent Coronary angiogram. Twelve patients underwent PTCA. One had Left main with Double vessel disease and was awaiting CABG.

Primary PTCA was possible in only 4 out of 69 patients. Majority had single vessel disease (61% of those who underwent CAG). LAD was the most common Culprit vessel seen in 8 out of 13 angiograms (61.5%).

This study is one of the largest among Indian studies on LVFWR in terms of number of subjects involved. There is a paucity of data from the Indian population regarding LVFWR. In this study we looked into the characteristics of AMI patients who died due to in-hospital LVFWR.

LVFWR following AMI usually occurs in elderly patients between 65 and 70 years of age.⁶⁻⁸ In our study, the mean age of subjects was about 65 years. Age group of 61-70 y constituted about 40% of subjects. Our study also showed a marginal female predominance which is consistent with previous data.⁹⁻¹¹

LVFWR is more often reported to occur after the first (transmural) myocardial infarction.^{8,12} In our study only eight out of 69 patients had past history of IHD. Diabetes mellitus and peripheral vascular disease are less incidental in these patients, as these conditions are associated with the development of collateral circulation, which may protect from cardiac rupture.¹² In our study Diabetes mellitus was seen in less than 30% of subjects.

Previous studies showed a higher incidence of arterial hypertension in LVFWR patients.^{7,13,14} One of the triggering factors for LVFWR is the presence of persistent arterial hypertension during the first 24 hours. Our study showed prevalence of hypertension in 38% of subjects.

Present study showed an obvious finding of a delay in hospital admission in the subjects. Mean duration of presentation to the hospital after symptom onset is around 23 hours. This correlated with previous data, which showed a delay in hospital admission as one of the major predictor of LVFWR.^{8,15} In an Indian study by Swaroop et al median duration of presentation to the hospital after the onset of symptoms was 24 h.¹⁶

In our study a high incidence of patients presenting in cardiogenic shock was observed (43%). A study by Swaroop et al¹⁶ showed similar high rate of cardiogenic shock upto 28.6% upon presentation. In the present analysis, 63.7% of patients had anterior wall STEMI. This correlated with the previous data which found anterior location of infarction has a higher incidence of LVFWR.

In some studies, administration of thrombolytics has been related to an increase in LVFWR,¹⁷⁻¹⁹ while in others, it was observed less frequently.^{20,21} Other than in a few trials, a multivariable analysis failed to identify the use of lytics as an independent risk factor for LVFWR. In our study, proportion of patients who received thrombolysis was 71% and primary PTCA was received by only 5.7% of patients. Late or failed thrombolysis is usually linked to increased rates of LVFWR.^{22,23} In our study, 5 patients had failed thrombolysis and all of them underwent successful rescue PTCA of culprit artery. PCI independently reduces the risk of LVFWR in comparison with thrombolysis.²⁴ Paradoxically, despite conferring an overall mortality-rate benefit, fibrinolytic agents have been implicated in the accelerated occurrence of LVFWR within the first 24 to 48 h after AMI.²⁵ Cardiac rupture is prevented by early thrombolysis therapy but is promoted by late treatment.

Previous data suggest that approximately 50% of patients with free-wall rupture are diagnosed within 5 days of the AMI, with 90% diagnosed within 2 weeks. In our study, 57.9% had LVFWR less than 48 h of AMI.

In the study by Swaroop et al.¹⁶ majority of the LVFWR patients presented with a single-vessel disease. Our study reinforced this, as 61.5% of subjects had SVD. A study by Markowicz et al reported that the most common culprit coronary lesion associated with LV rupture as the left anterior descending artery in 42% of patients, the circumflex coronary artery in 40% of patients, and the right coronary artery in 18% of patients.²⁶ In our study, out of 12 patients who underwent PTCA, 66.6% had LAD, 25% LCX and 8.3% RCA as the culprit coronary artery.

Angiographic reperfusion success is the most significant protective factor from FWR.¹ In our study 16 (23.1%) patients did not undergo any reperfusion as they were out of window period for lysis and had cardiac rupture before CAG was planned.

The treatment for LVFWR is mainly supportive in the immediate setting. None of the patient underwent surgery, as they did not survive till the surgery was undertaken. In our study usage of β -blockers among subjects was meager 21.7 % due to coexisting contraindications. Use of β -blockers and ACEI/ARB have all contributed to lower incidence and death rate of FWR as previously reported.^{27,28}

Limitations

1. This is a single-center- based, retrospective study
2. No patients who underwent surgery could be included in the study.
3. Only acute forms of LVFWR were selected
4. Number of patients who underwent primary PTCA was very less as compared to thrombolysis group.

Conclusion

This study reinforces the contributions of Classical predictors of LVFWR after myocardial infarction (AMI) in Indian setup, viz: advanced age, female gender, delayed presentation, presence of hypertension, first acute MI and cardiogenic shock at the time of presentation. Majority of LVFWR had single vessel disease predominantly LAD. β -blocker usage was considerably less in LVFWR patients. As majority of LVFWR patients had undergone thrombolysis, it is conceptualized that reduction of LVFWR may be achieved with PTCA in place of thrombolysis. This study underscores the fact that PTCA should be the modality of choice in the treatment of STEMI for prevention of complications like LVFWR.

References:

1. Nishiyama K, Shinichi O, Jirou A, et al. Coronary angioplasty reduces free wall rupture and improves mortality and morbidity of acute myocardial infarction. *J InvasCardiol* 2004;16:554–558.
2. Moreno R, Lopez-Sendon J, Garcia E, et al. Primary angioplasty reduces the risk of left ventricular free wall rupture compared with thrombolysis in patients with acute myocardial infarction. *J Am CollCardiol* 2002;39:598–603.
3. Nakamura F, Minamino T, Higashino Y, et al. Cardiac free wall rupture in acute myocardial infarction: ameliorative effect of coronary reperfusion. *ClinCardiol* 1992;15:244–250.
4. Raitt MH, Kraft CD, Gardner CJ, et al. Subacute ventricular free wall rupture complicating myocardial infarction. *Am Heart J* 1993;126:946–955
5. Mishra PK, Pathi V and Murday A. Post myocardial infarction free wall rupture. *Interact CardiovascThoracSurg* 2007; 6: 39–42.
6. London RE and London SB. Rupture of the heart. A critical analysis of 47 consecutive autopsy cases. *Circulation* 1965; 31: 202–208.
7. Naiem F, De la Maza LM and Robbins SL. Cardiac Rupture during myocardial infarction. A review of 44 cases. *Circulation* 1972; 45: 1231–1239.

8. Figueras J, Cortadellas J, Calvo F, Soler-Soler J. Relevance of delayed hospital admission on development of cardiac rupture during acute myocardial infarction: study in 225 patients with free wall, septal or papillary muscle rupture. *Journal of the American College of Cardiology*. 1998; 32: 135-139.
9. Mann JM, Roberts WC. Rupture of the left ventricular free wall during acute myocardial infarction: analysis of 138 necropsy patients and comparison with 50 necropsy patients with acute myocardial infarction without rupture. *The American journal of cardiology*. 1988; 62: 847-859.
10. Oliva PB, Hammill SC, Edwards WD. Cardiac rupture, a clinically predictable complication of acute myocardial infarction: Report of 70 cases with clinicopathologic correlation. *Journal of the American College of Cardiology*. 1993; 22: 720-726.
11. Purcaro A, Costantini C, Ciampani N, Mazzanti M, Silenzi C, Gili A, et al. Diagnostic criteria and management of subacute ventricular free wall rupture complicating acute myocardial infarction. *The American journal of cardiology*. 1997; 80: 397-405.
12. Friedman HS, Kuhn LA and Katz AM. Clinical and electrocardiographic features of cardiac rupture following acute myocardial infarction. *Am J Med* 1971; 50: 709-720.
13. Shapira I, Isakov A, Burke M, Almog C. Cardiac rupture in patients with acute myocardial infarction. *Chest*. 1987; 92: 219-223.
14. Wessler S, Zoll PM, Schlesinger MJ. The pathogenesis of spontaneous cardiac rupture. *Circulation*. 1952; 6: 334-351.
15. Figueras J, Cortadellas J, Soler-Soler J. Left ventricular free wall rupture: clinical presentation and management. *Heart* 2000; 83:499-504.
16. Varghese S, Ohlow M-A. Left ventricular free wall rupture in myocardial infarction: A retrospective analysis from a single tertiary center. *JRSM Cardiovasc Dis*. 2019;8: 2048004019896692.
17. Amery A, Roeber G, Vermeulen HJ, Verstraete M. Single-blind randomized multicentre trial comparing heparin and streptokinase treatment in recent myocardial infarction. *Acta Med Scand Suppl* 1969;505:1-35.
18. Heikinheimo R, Ahrenberg P, Honkapohja H, Iisalo E, Kallio V, Konttinen Y, Leskinen O, Mustaniemi H, Reinikainen M, Siitonen L. Fibrinolytic treatment in acute myocardial infarction. *Acta Med Scand* 1971;189:7-13.
19. Okino S, Nishiyama K, Ando K, Nobuyoshi M. Thrombolysis increases the risk of free wall rupture in patients with acute myocardial infarction undergoing percutaneous coronary intervention. *J Interv Cardiol* 2005;18:167-172.
20. European Working Party. Streptokinase in recent myocardial infarction: a controlled multicentre trial. *Br Med J* 1971; 3:325-331.
21. European Cooperative Study Group for Streptokinase Treatment in Acute Myocardial Infarction. Streptokinase in acute myocardial infarction. *N Engl J Med* 1979; 301:797-802.

22. Becker RC, Charlesworth A, Wilcox RG, Hampton J, Skene A, Gore JM, et al. Cardiac rupture associated with thrombolytic therapy: impact of time to treatment in the Late Assessment of Thrombolytic Efficacy [LATE] study. *Journal of the American College of Cardiology*. 1995; 25: 1063-1068.
23. Ohishi F, Hayasaki K, Honda T. Effect of thrombolysis on rupture of the left ventricular free wall following acute myocardial infarction. *Journal of Cardiology*. 1996; 28: 27-32.
24. Moreno R, Lopez-Sendon J, Garcia E, et al. Primary angioplasty reduces the risk of left ventricular free wall rupture compared with thrombolysis in patients with acute myocardial infarction. *J Am CollCardiol* 2002;39:598–603. Raitt MH, Kraft CD, Gardner CJ, et al. Subacute ventricular free wall rupture complicating myocardial infarction. *Am Heart J* 1993;126:946–955
25. Patel MR, Meine TJ, Lindblad L, et al. Cardiac tamponade in the fibrinolytic era: analysis of >100,000 patients with ST-segment elevation myocardial infarction. *Am Heart J* 2006; 151: 316–322.
26. Markowicz-Pawlus E, Nozynski J, Duszanska A, et al. The impact of a previous history of ischaemic episodes on the occurrence of left ventricular freewall rupture in the setting of myocardial infarction. *Kardiol Pol* 2012;70(7):713–7.
27. Figueras J, Alcalde O, Barrabés JA, et al. Changes in hospital mortality rates in 425 patients with acute ST-elevation myocardial infarction and cardiac rupture over a 30-year period. *Circulation* 2008;118:2783–9.
28. Sulzgruber P, El-Hamid F, Koller L, et al. Long-term outcome and risk prediction in patients suffering acute myocardial infarction complicated by post-infarction cardiac rupture. *Int J Cardiol* 2017;227:399–403.