

ASSESSMENT QUALITY OF LIFE IN EGYPTIAN CHILDREN AND ADOLESCENTS WITH EPILEPSY

Safaa Hamdy Ahmad Saleh ⁽¹⁾, Mohammed Elsayed Hamed Mohammed ⁽¹⁾ and *Osamah Abdullah Abounaji ⁽²⁾

⁽¹⁾ Professor of Pediatrics, Faculty of Medicine – Zagazig University

⁽²⁾ M.B.B.Ch, Faculty of Medicine – Elmergib University – Libya, Physician Visitor to Zagazig University Children Hospital

* Corresponding author : Osamah Abdullah Abounaji
e-mail: aosamah32@yahoo.com

Abstract

Background: Epilepsy is a common chronic neurological condition in developing years that can negatively impact one's physical, social and emotional function. The aim of the study was to explore the current status of health related quality of life (HR-QOL) in Egyptian children and adolescents with epilepsy using (WHOQOL-BREF) questionnaire

Patients and Methods: This Case control study was carried out in pediatric neurology unit and outpatient clinic in department of pediatrics at Zagazig University Children's Hospital on 110 children. All the children were subjected to history taking, clinical examination and assessment of QOL by WHOQOL-BREF.

Results: There was a significant decreases in all domains and total score of WHOQOL-Bref questionnaire between studied group. There was a significant –ve correlation between age and psychological domain score, also between number of school failure and both psychological and environmental domain. Disease duration and treatment duration were –ve significantly correlated with all QOL domains. A significant +ve correlation was founded between social class score and both general and environmental domain.. There was a significance decrease in general and environmental domain score between cases with generalized and partial seizures compared to cases with absent seizures.

Conclusion: Diminished QOL is common in children with epilepsy than other children. Patients with epilepsy had lower mean scores of all domains of QOL, especially those with frequent fits, those with long duration, and in patients with generalized fits.

Keywords: Quality of life – Childhood – Epilepsy – Health

INTRODUCTION

Epilepsy has a significant emotional impact on parents of affected children, and parental emotional stability has been found to be a major predictor of the quality of life (QOL) of the epileptic child. It has been associated with significant psychosocial maladjustment in both the affected children and their families ⁽¹⁾.

Childhood epilepsy is an elevated hazard for poor psychological outcomes and impacts on quality of life of children but also has a great sway on family functioning. Children being the high-risk group and in crucial development period during which many cognitive and social skills have to be learned QOL is a significant health outcome to assess children with epilepsy ⁽²⁾.

Epilepsy also has a significant emotional impact on parents of affected children, and parental emotional stability has been found to be a major predictor of the quality of life (QOL) of the epileptic child. It has been associated with significant psychosocial maladjustment in both the affected children and their families ⁽³⁾.

Children with epilepsy experience difficulty in aspects of functioning, including emotional and behavioral problems, Social competence, academic achievement, and family life, with effects extending into adulthood. The management of epilepsy requires recognition of potential effects of epilepsy and all aspects of life ⁽³⁾.

Quality of life (QOL) is affected by age, seizure frequency, parent's education, type of epilepsy, and type of anti epileptic in children with epilepsy. Cognition, energy levels and concentration are most commonly affected due to epilepsy ⁽⁵⁾.

The aim of the study was to explore the current status of health related quality of life (HR-QOL) in Egyptian children and adolescents with epilepsy by using (WHOQOL-BREF) questionnaire.

PATIENTS AND METHODS

This Case control study was carried out in pediatric neurology unit and outpatient clinic in department of pediatrics at Zagazig University Children's Hospital on 110 subjects, divided into two groups, case group included 55 children with epilepsy and control group included 55 apparently healthy children without epilepsy of matched age, sex and social class during the period from February 2021 to September 2021, to explore the current status of health related quality of life (HR-QOL) in Egyptian children and adolescents with epilepsy by using (WHOQOL-BREF) questionnaire and the Socioeconomic position (SEP) scale ⁽⁶⁾

Written informed consent was obtained from all children' parents or their relatives and the study was approved by the research ethical committee of Faculty of Medicine, Zagazig University. The work has been

carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for studies involving humans

Inclusion Criteria:

Enrollment criteria included:

1. Aged between 9 and 25 years,
2. Both Sex,
3. Ability to answer the questions by himself/herself,
4. Medical diagnosis of epilepsy for more than 2 years.

Exclusion criteria:

The exclusion criteria for both groups included:

1. Children < 9 and > 16 years
2. Previous brain surgery,
3. Use of a concomitant medication with central nervous system effects, or presence of another progressive neurological , psychiatric illness or other chronic disease.

All patients and controls will be subjected to the following:

- History taking obtained for each child included age, sex, child's hobbies and leisure time activities, sleep disorders, urine incontinence, previous academic failure or success, and last year's achievements and absenteeism rate.
- Epilepsy was defined as separate occurrence of two or more unprovoked seizures, not diagnosed as neonatal or febrile seizures.
- The diagnosis of epilepsy in the children based on history and the electroencephalographic findings, as all of them underwent electroencephalography.

Clinical examination :

- All the studied children were subjected to full clinical examination to exclude other medical conditions and chronic diseases.

Assessment of HR-QOL of all the studied children and adolescents It was held by using WHOQOL-BREF questionnaire. ⁽⁷⁾

The questionnaire consisted of 26 items but one item about appreciation of sexual life was discarded putting into consideration the young age of the studied group. The items assessing four main domains that cover the aspects proposed to judge QOL.

The items assessing four main domains that cover the aspects proposed to judge QOL. The assessed domains had included : Physical health, Psychological health, Social relationships and Environmental domain.

Data collection

Data were collected by using an interview questionnaire that included the following updated social scale:

- (1) Mother's education.
- (2) Father's education.
- (3) Working status of the mother.
- (4) Working status of the father.
- (5) Use of computer.
- (6) Per-capita income.
- (7) Family size.
- (8) Crowding index.
- (9) Proper sewage disposal.
- (10) Proper refuse disposal.

The following changes were made to the modified scale :

- (1) Working status was separated from education of both the father and the mother.
- (2) A new educational domain, the use of computer, was added.
- (3) Home sanitation included only proper sewage and refuse disposal as water supply was found to be available for all the families studied.

Statistical analysis

Data collected throughout history, basic clinical examination, laboratory investigations and outcome measures coded, entered and analyzed using Microsoft Excel software. Data were then imported into Statistical Package for the Social Sciences (SPSS version 20.0) (Statistical Package for the Social Sciences) software for analysis. According to the type of data qualitative represent as number and percentage, quantitative continues group represent by mean \pm SD, the following tests were used to test differences for significance; .difference and association of qualitative variable by Chi square test (X^2). Differences between quantitative independent groups by t test. P value was set at <0.05 for significant results and <0.001 for high significant result.

RESULTS

Table (1): Demographic data of the studied groups:

Variable		Cases (n=55)		Control (n=55)		t	P
Age: (years)	Mean ± Sd	12.69±2.68		12.33±2.4		0.75	0.46
	Range	9 – 16		9 - 16			
Variable		No	%	No	%	χ ²	P
Sex:	Female	22	40	23	41.8	0.04	0.85
	male	33	60	32	58.2		
Mother education:	Illiterate	11	20	8	14.5	8.7	0.12
	Read & write	5	9.1	3	5.5		
	Primary	2	3.6	11	20		
	Preparatory	9	16.4	7	12.7		
	Secondary	9	16.4	5	9.1		
University	19	34.5	21	38.2			
Father education:	Illiterate	5	9.1	1	1.8	6.55	0.26
	Read & write	3	5.5	2	3.6		
	Primary	4	7.3	1	1.8		
	Preparatory	7	12.7	9	16.4		
	Secondary	8	14.5	14	25.5		
University	28	50.9	28	50.9			
Mother work:	Not working	35	63.6	31	56.4	0.61	0.44
	Working	20	36.4	24	43.6		
Father work:	Not working	5	9.1	3	5.5	0.54	0.46
	Working	50	90.9	52	94.5		
Social class:	Low	18	32.7	11	20	3.22	0.20
	Moderate	31	56.4	33	60		
	High	6	10.9	11	20		

SD: Standard deviation t: Independent t test χ²:Chi square test NS: Non significant (P>0.05)

Table (1) showed there were no statistical significant differences between the studied groups regarding demographic data.

Table (2): Epilepsy data among the studied cases group:

Variable		Cases (n=55)	
Age of onset: (years)	Mean ± Sd	6.24±3.45	
	Range	2-13	
	Median (IQR)	5(3-10)	
Duration: (years)	Mean ± Sd	6.45±2.93	
	Range	3-14	
	Median (IQR)	6(4-8)	
Variable		No	%
Type of epilepsy	Absence seizers	4	7.3
	Generalized tonic clonic	41	74.5
	Partial	10	18.2
Description of attacks	Convulsion	50	90.9
	Relaxation with loss of consciousness	5	9.1
Number of attack/month	1-2	47	87.3
	3-4	5	9.1
	5-7	2	3.6
Attacks at school	No	40	72.7
	Yes	15	27.3
Headache before attack	No	19	34.5
	Yes	36	65.5
Psychological tension before attack	No	18	32.7
	Yes	37	67.3
Fever before attack	No	45	81.8
	Yes	10	18.2
Exposure to bright light cause the attack	No	50	90.9
	Yes	5	9.1

SD: Standard deviation (IQR): Inter quartile range

Table (2) showed that mean age of onset among the cases group was 6.24 years while mean disease duration was 6.45 years. Most frequent type was generalized tonic clonic (74.5%) and most frequent attacks were convulsions. 87.3% of cases had from 1 to 2 attacks per month and 27.3% had attacks at school. Frequency of headache before attacks was 65.5%, psychological tension 67.3% and fever 18.2%. Finally in 9.1% of cases exposure to bright light cause the attack.

Table (3): WHOQOL-Bref questionnaire results among the studied groups:

Domain		Cases (n=55)	Control (n=55)	t	P
General health	Mean ± Sd	44.77±12.66	75.45±9.31	14.48	<0.001**
	Range	25-100	62.5-100		
Physical health	Mean ± Sd	42.92±14.49	84.48±7.78	18.74	<0.001**
	Range	17.86-92.86	71.43-100		
Psychological health	Mean ± Sd	44.77±11.31	69.32±7.41	13.46	<0.001**
	Range	20.83-87.5	50-87.5		
Social life	Mean ± Sd	48.18±12.13	60.68±8.13	6.35	<0.001**
	Range	25-100	50-75		
Environmental	Mean ± Sd	45.28±10.04	64.43±8.01	11.05	<0.001**
	Range	25-93.75	31.25-78.13		
Total	Mean ± Sd	35.75±7.97	57.44±4.62	17.46	<0.001**
	Range	23.2-74.4	47.2-68		

SD: Standard deviation t: Independent t test **: highly significant (P<0.001)

Table (3) showed that there was a statistical significant decreases in all domains and total score of WHOQOL-Bref questionnaire among cases group compared to control group.

Table (4): Correlation between WHO-QOL questioner domain and some parameters among the studied cases group:

Variable	General (n=55)		Physical (n=55)		Psychological (n=55)		Social (n=55)		Environment (n=55)	
	r	P	r	P	r	P	r	P	R	P
Age (years)	-0.03	0.84 NS	-0.02	0.92 NS	-0.25	0.03*	0.02	0.90 NS	-0.10	0.48 NS
Social class score	0.31	0.02*	0.09	0.52 NS	0.03	0.84 NS	-0.05	0.73 NS	0.37	0.009*
N school failure	-0.14	0.14 NS	0.12	0.40 NS	-0.26	0.03*	0.04	0.77 NS	-0.33	0.01*
Age of onset(y)	0.19	0.17 NS	0.09	0.52 NS	-0.05	0.71 NS	0.20	0.06 NS	0.18	0.12 NS
Duration(y)	-0.26	0.03*	-0.25	0.04*	-0.27	0.02*	-0.25	0.04*	-0.34	0.01*
Attack/m	0.02	0.90 NS	-0.26	0.03*	-0.34	0.01*	-0.27	0.03*	0.05	0.75 NS
Duration of ttt(y)	-0.27	0.03*	-0.28	0.03*	-0.26	0.03*	-0.26	0.03*	-0.32	0.01*
N. of drugs	0.07	0.61 NS	-0.04	0.78 NS	-0.11	0.42 NS	-0.16	0.24 NS	-.008	0.56 NS
N. hospitalization	-0.03	0.87 NS	0.01	0.92 NS	0.07	0.63 NS	0.08	0.58 NS	0.02	0.88 NS

r: Pearson's correlation coefficient NS: Non significant (P>0.05) *: Significant (P<0.05)

Table (4) showed there was a statistical significant -ve correlation between age and psychological domain score also between number of school failure and both psychological and environmental domain. Disease duration and treatment duration were -ve significantly correlated with all QOL domains. A statistical significant +ve correlation was founded between social class score and both general and environmental domain.

Table (5): Relation between WHO-QOL questioner domain and Type of epilepsy among the studied cases group:

Domain	Type	N	Mean	SD	F	P
General	Absence seizers	4	59.38	27.72	3.18	0.03*
	Generalized tonic	41	43.29	10.86		
	Partial	10	45.00	8.74		
Physical	Absence seizers	4	52.68	27.87	1.07	0.35 NS
	Generalized tonic	41	41.72	13.93		
	Partial	10	43.93	9.38		
Psychological	Absence seizers	4	52.08	23.69	0.91	0.41 NS
	Generalized tonic	41	44.31	10.53		
	Partial	10	43.75	7.92		
Social	Absence seizers	4	59.38	27.72	1.9	0.16 NS
	Generalized tonic	41	47.26	10.27		
	Partial	10	47.50	9.86		
Environmental	Absence seizers	4	56.25	25.13	3.01	0.04*
	Generalized tonic	41	44.89	8.02		
	Partial	10	42.50	6.78		

SD: Standard deviation t: Independent t test NS: Non significant (P>0.05) *: Significant(p<0.05)

Table (6) showed that there was a statistical significance decrease in general and environmental domain score among cases that had generalized and partial seizers compared to cases that had absent seizers. Physical, psychological and social domain also were decreased in cases that had generalized and partial seizers compared to cases that had absent seizers but without statistical significance

DISCUSSION

Epilepsy is a common chronic neurological condition in developing years that can negatively impact one’s physical, social and emotional function. In Austria reported behavioral and emotional problems in 22% of epileptic children, and studies have described feelings of shame, rejection, fear, worry, low self esteem, and perception of stigma to be common in children with epilepsy⁽⁸⁾.

Increasing attention is being focused on problems experienced by children with epilepsy as a result of stigma, which is associated with poor psychosocial health outcomes and impaired QOL. Epilepsy may interfere with social functioning by limiting educational opportunities, employability, and interpersonal relationships and also increase the risk of death⁽⁹⁾.

Our study shows that there were no statistical significant differences between the studied groups regarding demographic data.

Monir et al. ⁽¹⁰⁾ found that there were no statistical significant differences between the studied groups regarding demographic data.

Sherman et al. ⁽¹¹⁾ found that sociodemographic factors were found to be weak predictors of a poorer total QOL score in children with epilepsy.

Our study cleared that mean age of onset among the cases group was 6.24 years while mean disease duration was 6.45 years. Most frequent type was generalized tonic clonic (74.5%) and most frequent attacks were convulsions. 87.3% of cases had from 1 to 2 attacks per month and 27.3% had attacks at school. Frequency of headache before attacks was 65.5%, psychological tension 67.3% and fever 18.2%. Finally in 9.1% of cases exposure to bright light cause the attack.

Monir et al. ⁽¹⁰⁾ showed a comparison of the types of seizures and epilepsy-related variables in the studied patient groups. Highly significant differences (P < 0.001) were found between patient subgroups regarding the age at onset of seizures, the duration of therapy, the number of antiepileptic drugs, and seizure frequency per month.

Our study reported that mean treatment duration among the cases group was 6.17 years. Almost three quarters of the cases take 1 AED. Also, 70.9% were adherent to drug therapy and 69.1% control the attacks for weeks, 73.6% of the cases admitted to hospital because of epilepsy. Our study illustrated that there was a statistaical significant decreases in all domains and total score of WHOQOL-Bref questionnaire among cases group compared to control group.

Miller et al. ⁽¹²⁾, who demonstrated that the HRQOL in children with epilepsy compared with healthy controls is diminished in all functioning domains

Our study revealed that there was a statistical significant –ve correlation between age and psychological domain score also between number of school failure and both psychological and environmental domain. Disease

duration and treatment duration were –ve significantly correlated with all QOL domains. A statistical significant +ve correlation was founded between social class score and both general and environmental domain.

Abbas et al. ⁽¹³⁾ was found to have a significantly poor QOL in children with epilepsy and longer seizure and treatment durations.

Guekht et al. ⁽¹⁴⁾ reported that patients with frequent seizures had a lower QOL score, which is in agreement with our results.

Benavente et al. ⁽¹⁵⁾ who also found that Duration of the illness, severity of epilepsy and seizure frequency have been found to be related to diminished QOL in children with epilepsy .

Our study revealed that there was no statistical significant differences between presence or absence of nocturnal enuresis and QOL domain among studied cases group. Our study revealed that there was a statistical significance decrease in general and environmental domain score among cases that had generalized and partial seizers compared to cases that had absent seizers. Physical, psychological and social domain also were decreased in cases that had generalized and partial seizers compared to cases that had absent seizers but without statistical significance.

Monir et al. ⁽¹⁰⁾ showed that seizures control were found to be significantly strong predictors of an impaired general and environmental domain score.

Conclusion: Diminished QOL is common in children with epilepsy. Patients with epilepsy had lower mean scores of all domains of QOL, especially those with frequent fits, those with long duration, and in patients with generalized fits.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request

Conflicting Interest (If present, give more details): No Conflict of Interest

No financial disclosure

-Acknowledgements

Not applicable

Declarations

-Ethics approval and consent to participate

Written informed consent was obtained from all patients and the study was approved by the research ethical committee of Faculty of Medicine, Zagazig University (International review board ZU-IRB #6340/30-1-2021). The study was done according to The Code of Ethics of the World Medical Association (Declaration of Helsinki) for studies involving humans.

-Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

REFERENCES

- 1- Aronu, A. E., Uwaezuoke, N., Chinawa, J. M., Bisi-Onyemaechi, A., & Ojinnaka, N. C. (2021).** Health-related quality of life in children and adolescents with epilepsy in Enugu: Need for targeted intervention. *Nigerian Journal of Clinical Practice*, 24(4), 517.
- 2- Taylor, J., Jacoby, A., Baker, G. A., & Marson, A. G. (2011).** Self reported and parent - reported quality of life of children and adolescents with new onset epilepsy. *Epilepsia*, 52(8), 1489-1498.
- 3- Nawani, H., Smith, M. L., Wheeler, A. L., & Widjaja, E. (2019).** Functional Connectivity Associated with Health-Related Quality of Life in Children with Focal Epilepsy. *American Journal of Neuroradiology*, 40(7), 1213-1220.
- 4- Karnavat, P. K., Hegde, A. U., & Kulkarni, S. (2018).** Quality of life in children with epilepsy in private and public tertiary care centers in India. *International Journal of Epilepsy*, 5(01), 028-037.
- 5- Nagarathnam, M., Shalini, B., Vijayalakshmi, V., Vengamma, B., & Latheef, S. A. A. (2017).** Predictors of quality of life among adolescents with epilepsy in the state of Andhra Pradesh. *Neurology India*, 65(5), 1019.
- 6- Fahmy, S. I., Nofal, L. M., Shehata, S. F., El Kady, H. M., & Ibrahim, H. K. (2015).** Updating indicators for scaling the socioeconomic level of families for health research. *Journal of the Egyptian Public Health Association*, 90(1), 1-7.
- 7- Golami A, Jahormi L, Zari E, Dehghan A. (2013).** The world health organization's WHOQOL-BREF quality of life assessment: psychometric properties and results of the international field trial. A report from the WHOQOL group .*int j Prev Med.*; 4(7): 809-817
- 8- Vuong, E., Nothling, J., Lombard, C., Jewkes, R., Peer, N., Abrahams, N., & Seedat, S. (2020).** Peripheral adiponectin levels in anxiety, mood, trauma-and stressor-related disorders: A systematic review and meta-analysis. *Journal of affective disorders*, 260, 372-409.

- 9- **Yogarajah, M., & Mula, M. (2019).** Social cognition, psychiatric comorbidities, and quality of life in adults with epilepsy. *Epilepsy & Behavior*, 100, 106321.
- 10- **Monir, Zeinab & El-Alameey, Inas & Eltahlawy, Eman. (2013).** Health related quality of life of children with epilepsy in Egypt. *Journal of The Arab Society for Medical Research*. 8. 53. 10.4103/1687-4293.123787.
- 11- **Sherman EMS, Griffiths SY, Akdag S, Connolly MB, Slick DJ, Wiebe S. (2008).** Sociodemographic correlates of health-related quality of life in pediatric epilepsy. *Epilepsy Behav* 2008; 12:96-101. †
- 12- **Miller V, Palermo TM, Grewe S. (2001).** Health-related quality of life in children with epilepsy 2001; Gainesville, FL: Presented at the 8th Florida Conference on Child Health Psychology
- 13- **Abbas, Zahia, Maha A. Elseed, and Inaam N. Mohammed. (2014).** The quality of life among Sudanese children with epilepsy and their care givers." *Sudanese journal of paediatrics* 14.1: 51.
- 14- **Guekht AB, Mitrokhina TV, Lebedeva AV, Dzugaeva FK, Milchakova LE, Lokshina OB, et al. (2007).** Factors influencing on quality of life in people with epilepsy. *Seizure*; 16:128–133.
- 15- **Benavente-Aguilar I, Morales-Blázquez C, Rubio EA, Rey JM. (2004).** Quality of life of adolescents suffering from epilepsy living in the community. *J Paediatr Child Health*; 40:110-113.