

# IMPLEMENTATION OF ENHANCED RECOVERY AFTER SURGERY (ERAS) PROTOCOLS IN ELECTIVE GENERAL SURGERIES AT A PERIPHERAL SECONDARY CARE HOSPITAL IN PUNJAB: A PROSPECTIVE OBSERVATIONAL STUDY

**DR. GURSIMRAN SINGH**

**DR. SUKHDEEP KAUR**

**DR. ASISH GUPTA**

**DR. NAVKIRAT SINGH**

Sub Divisional Hospital (SDH), Patti, District Tarn Taran, Punjab, India

Corresponding Author: Dr. Gursimran Singh

Email: [dr.gursimran@yahoo.co.in](mailto:dr.gursimran@yahoo.co.in)

Phone: 919814031400

## ABSTRACT

Enhanced Recovery After Surgery (ERAS) is a multimodal perioperative care pathway aimed at reducing surgical stress and improving postoperative recovery. This prospective observational study evaluated the feasibility and outcomes of ERAS protocols implementation in elective general surgeries at a peripheral secondary care hospital in Punjab. One hundred adult patients undergoing elective general surgical procedures were enrolled over a period of 10 months. ERAS measures included reduced fasting duration, carbohydrate loading, multimodal analgesia, early oral feeding, and early mobilization. The mean age of patients was  $42.6 \pm 12.4$  years. Open cholecystectomy constituted 69% of procedures, followed by hernia surgery and exploratory laparotomy. Early oral intake within 24 hours was achieved in 98% of patients, while mobilization within 24 hours was achieved in all patients. Mean hospital stay was  $3.2 \pm 1$  days. No mortality or major complications occurred. ERAS implementation was found to be feasible, safe, and beneficial in a resource-limited peripheral hospital setting.

## KEYWORDS

ERAS; General Surgery; Peripheral Hospital; Enhanced Recovery

## INTRODUCTION

Enhanced Recovery After Surgery (ERAS) protocols are evidence-based perioperative pathways designed to reduce surgical stress and improve postoperative outcomes. ERAS pathways have shown benefits including reduced hospital stay, lower complication rates, and faster patient recovery in multiple surgical specialties [1-3]. Despite increasing adoption in tertiary healthcare institutions, implementation in peripheral secondary care hospitals remains limited due to infrastructural, financial, and manpower constraints. This study was conducted to evaluate the feasibility, safety, and outcomes associated with implementation of ERAS protocols in elective general surgeries at a peripheral secondary care hospital in Punjab.

## **MATERIALS AND METHODS**

This prospective observational cohort study was conducted in the Department of General Surgery at Sub Divisional Hospital Patti, Punjab, India, between June 2025 and March 2026 . Adult patients aged 18–70 years undergoing elective general surgical procedures were included after obtaining written informed consent. Patients undergoing emergency surgery and those with ASA grade IV or higher were excluded.

ERAS measures implemented included preoperative counseling, reduced fasting duration, carbohydrate loading, multimodal analgesia, early oral feeding, and early mobilization. Outcome measures included hospital stay duration, time to oral intake, mobilization time, and postoperative complications.

Data analysis was performed using Microsoft Excel 2021. Continuous variables were expressed as mean  $\pm$  standard deviation, while categorical variables were presented as frequencies and percentages.

## **ETHICAL CONSIDERATIONS**

The study protocol was approved by the Institutional Ethics Committee of Sub Divisional Hospital Patti. Written informed consent was obtained from all participants. The study was conducted in accordance with the Declaration of Helsinki ethical principles.

## **RESULTS**

A total of 108 patients were screened for eligibility. Eight patients were excluded due to emergency surgery or ASA grade IV status. Finally, 100 patients were included in the analysis.

The mean age was  $42.6 \pm 12.4$  years. Open cholecystectomy was performed in 69 patients (69%), hernia surgery in 21 patients (21%), and exploratory laparotomy in 10 patients (10%). Early oral intake within 24 hours was achieved in 98 patients (98%), while mobilization within 24 hours was achieved in all patients (100%). Mean hospital stay was  $3.2 \pm 1$  days. Postoperative complications occurred in 2 patients (2%) and consisted only of postoperative nausea and vomiting. No mortality or major complications were observed.

## **DISCUSSION**

The present study demonstrates that ERAS protocols implementation in a peripheral secondary care hospital is feasible and associated with favorable postoperative outcomes. Early oral intake and mobilization were successfully achieved in the majority of patients.

The low postoperative complication rate further supports the safety of ERAS pathways in resource-limited settings.

This study contributes valuable evidence regarding ERAS protocols implementation in rural healthcare environments and highlights the practicality of simplified ERAS pathways in peripheral government hospitals.

Limitations of the study include the observational design, absence of a control group, relatively small sample size, and limited follow-up duration.

## **CONCLUSION**

Implementation of ERAS protocols in elective general surgeries at a peripheral secondary care hospital in Punjab is feasible, safe, and beneficial. Structured perioperative pathways and staff training may further improve postoperative outcomes in resource-limited healthcare settings.

## **ACKNOWLEDGEMENT**

The authors acknowledge the surgical staff and patients of Sub Divisional Hospital Patti for their support and participation in this study.

## **FUNDING**

The authors received no financial support for the research, authorship, or publication of this article.

## **CONFLICT OF INTEREST**

The authors declare no conflict of interest.

## **REFERENCES**

1. Kehlet H, Wilmore DW. Multimodal strategies to improve surgical outcome. *Am J Surg.* 2002;183(6):630-641.
2. Ljungqvist O, Scott M, Fearon KC. Enhanced Recovery After Surgery: A review. *JAMA Surg.* 2017;152(3):292-298.
3. Fearon KC, Ljungqvist O, Von Meyenfeldt M, et al. Enhanced recovery after surgery: A consensus review of clinical care. *Clin Nutr.* 2005;24(3):466-477.

4. Gustafsson UO, Scott MJ, Hubner M, et al. Guidelines for perioperative care in elective colorectal surgery: Enhanced Recovery After Surgery Society recommendations. *World J Surg.* 2019;43(3):659-695.
5. Thiele RH, Rea KM, Turrentine FE, et al. Standardization of care: impact of an enhanced recovery protocol on length of stay, complications, and direct costs after colorectal surgery. *J Am Coll Surg.* 2015;220(4):430-443.
6. Varadhan KK, Neal KR, Dejong CH, Fearon KC, Ljungqvist O, Lobo DN. The enhanced recovery after surgery pathway for patients undergoing major elective open colorectal surgery: A meta-analysis. *Clin Nutr.* 2010;29(4):434-440.
7. Nicholson A, Lowe MC, Parker J, Lewis SR, Alderson P, Smith AF. Systematic review and meta-analysis of enhanced recovery programmes in surgical patients. *Br J Surg.* 2014;101(3):172-188.
8. Miller TE, Roche AM, Mythen M. Fluid management and goal-directed therapy as an adjunct to Enhanced Recovery After Surgery (ERAS). *Can J Anaesth.* 2015;62(2):158-168.
9. Ljungqvist O. ERAS—Enhanced Recovery After Surgery: Moving evidence-based perioperative care to practice. *JPEN J Parenter Enteral Nutr.* 2014;38(5):559-566.
10. Nelson G, Bakkum-Gamez J, Kalogera E, et al. Guidelines for perioperative care in gynecologic/oncology surgery: Enhanced Recovery After Surgery Society recommendations. *Gynecol Oncol.* 2016;140(2):313-322.

**Table 1: Demographic Characteristics**

Variable	Value
Mean Age	42.6 ± 12.4 years
Male Patients	58
Female Patients	42
Sample Size	100
Study Duration	10 Months

**Table 2: Distribution of Surgical Procedures**

Procedure	Frequency (%)
Open Cholecystectomy	69 (69%)
Hernia Surgery	21 (21%)
Exploratory Laparotomy	10 (10%)

**Table 3: ERAS Outcome Parameters**

Outcome Parameter	Result	Percentage
Early Oral Intake (<24 h)	98 Patients	98%
Early Mobilization (<24 h)	100 Patients	100%
Postoperative Complications	2 Patients	2%
Mortality	0 Patients	0%