

EVALUATION OF ANTIBIOTIC PROPHYLAXIS USAGE IN SURGERY WARDS BEFORE AND AFTER THE IMPLEMENTATION OF CLINICAL GUIDELINES IN A TERTIARY CARE HOSPITAL- A RETROSPECTIVE STUDY

Dr Pandey¹ , Dr. S.Sathyavani²

1 Professor, Department of General Surgery, Sree Mookambika Institute of Medical Sciences Kanyakumari, Tamil Nadu, India.

2., Junior Resident, Department of General Surgery Sree Mookambika Institute of Medical Sciences College Kanyakumari, Tamil Nadu, India.

Corresponding Author S.Sathyavani ,Junior Resident, Department of General Surgery ,Sree Mookambika Institute of Medical Sciences College Kanyakumari, Tamil Nadu, India.

ABSTRACT :

Background: Wound infections are the commonest hospital-acquired infections in surgical patients. Approximately 30-50% of antibiotic use in hospitals, is now for surgical prophylaxis. However, between 30-90% of this prophylaxis is inappropriate, that increases the selective pressure favoring the emergence of antimicrobial resistance. Judicious use of antibiotics in the hospital through effective antibiotic policy and guideline development is thus essential.

Aims: To prepare a guideline for surgical antibiotic prophylaxis and to evaluate the impact of it in clinical practice.

Materials and Methods: An antibiogram was prepared to evaluate the sensitivity pattern retrospectively. A prospective study was done to evaluate the prescription pattern of antibiotic prophylaxis and incidence of surgical site infection. A draft guideline was prepared with expertise by the surgeon. Standardized recommendations in the guideline were done by the nominal group technique. Post-recommendation evaluation was done to observe the impact of the guideline in clinical practice.

Results and Conclusions: 222 cases were evaluated during the pre-recommendation period, of which 21 were infected with surgical site infections that were rated as 9.45%. Use of prophylactic antibiotics was widespread and inconsistent with published guidelines. 56 cases were evaluated during the post-recommendation period. None of the cases were infected. Prophylactic antibiotic use was consistent with respect to choice and duration. The study showed the importance of a multidisciplinary approach in implementing local guidelines in healthcare institutions, as per their hospital antimicrobial sensitivity pattern and expert's opinions.

Keywords: Antibiotic prophylaxis, guideline adherence or practice guidelines, guideline, humans

INTRODUCTION:

Surgical site infections (SSIs) are the second most common cause of nosocomial infections.[1] Though the post- Listerian era is enriched with much advances in field of asepsis and surgical and antiseptic techniques, SSIs are still a threat to the surgical world. Antibiotic prophylaxis is one component of preventive strategy against SSIs, that is based on a good surgical technique, strict asepsis in the operating theatre and control of infection within the hospital or general practice.[2],[3] Although properly administered antibiotics can reduce postoperative SSIs secondary to preoperative bacterial contamination, widespread use of prophylactic antibiotics have the disadvantages of emergence of multi-resistant organisms.[4]

The rich tropical climate of India additionally challenges the clinicians to manage a wide variety of infectious diseases. The higher rates of surgical site infections are associated not only with a higher morbidity and mortality, but also with increased medical costs. Hence, a constant awareness of the ever-present threat of infection must be a way of life for the entire surgical fraternity.

AIM AND OBJECTIVES OF THE STUDY:

To identify the common pathogens causing infections and their sensitivity pattern, to determine the incidence of surgical site infection, to prepare guidelines for the use of surgical antibiotic prophylaxis and to evaluate the impact of the same in clinical practice.

MATERIALS AND METHODS:

The study was conducted in the department of surgery, of a multidisciplinary teaching hospital of South tamilnadu, after being approved by the ethical committee. Initially a retrospective analysis of microbiological data (from medical records) of operated cases was analyzed from September 2023 to August 2024, to evaluate the susceptibility pattern of antibiotics. Subsequent follow- up chart reviews were also done to detect the incidence of SSIs. Each follow-up records were evaluated to determine the surgical site infections, that included pharmacy, microbiology, biochemistry and other lab records .

Then a prospective study was carried out over an eight month period, from September 2023 to April 2024. Inpatients undergoing surgical procedures under clean and clean-contaminated category were included and those having immunocompromised conditions, patients with active infection requiring treatment before or at the time of surgery and surgical procedures under contaminated and dirty category, were excluded. An antibiogram was prepared from the microbiology ledger to identify the common pathogens involved in the surgical infections, along with sensitivity pattern. A retrospective analysis of one-year data from September 2023 till September 2024 was done.

The prospective study involved data collection from patient case records and microbiology reports for demographic details and clinical details, including prophylactic antibiotic dose, duration, frequency and route. For detection of surgical site infections, the wound was inspected for any evidence of infection starting from 48 hours after surgery, to the day when the patient was discharged. Patients who

underwent laparoscopic procedures were discharged on the 3rd and 4th postoperative day. The rest of the patients were discharged after the suture removal, with clinically no evidence of SSIs.

The patients were called for follow-up after 1-2 weeks in case of laparoscopic cholecystectomy for the suture removal and after 15 days to a month in open cholecystectomy cases. Patients were observed during their follow-up visits in such cases. Self-reporting of infections by the patients was also encouraged, once they were discharged. The criterion for SSI was based on centre for disease control's (CDC's) definition of SSI. The patient who satisfied the superficial incisional surgical site infection criteria was studied. Cases were followed to know the exact time of administration of the prophylactic antibiotic in the operation theatre. The prescription pattern was evaluated with respect to choice, duration, route and frequency.

Duration of hospital stay was evaluated in the infected group and non-infected group. Data collected from the study, along with evidence-based literature and microbiology reports, were clubbed together to make a draft guideline, with expertise from a surgeon. This draft guideline was circulated among the surgical staff for their feedback on the recommendations made regarding the choice and duration of antibiotic prophylaxis in surgical procedures belonging to the clean and clean- contaminated surgery. Standardization of the draft guideline was done by a nominal group technique, where each recommendation was stated, followed by the feedback results and then a discussion was initiated among the surgeons to accomplish the standardization procedure. After standardization of the guideline, a 7-15 days interval was given and then the cases were studied to evaluate the change (if any) in utilization of surgical antibiotic prophylaxis.

Statistical analysis was done using the statistical package for social sciences (SPSS). Different statistical methods were used as appropriate. Mean \pm SD was determined for quantitative data and frequency for categorical variables. The independent t- test was performed on all continuous variables. The normal distribution data was checked before any t-test. The Chi-Square test was used to analyze group difference for categorical variables. A p- value < 0.05 was considered significant

RESULTS:

168 cases were evaluated to prepare an antibiogram retrospectively. *Escherichia coli* was the most common pathogen found [42 (25%)], followed by *Pseudomonas aeruginosa* [35 (20.83%)] and coagulase-positive staphylococci [32 (19.04%)] in surgical set-up. Of the antimicrobials commonly used, impressive results of susceptibility pattern were observed with an aminoglycoside, amikacin (61.30%), followed by a third generation cephalosporin, cefotaxime (40.47%) and amoxicillin-clavulanic acid (33.92%).

The prospective study evaluated a total of 308 cases, of which 222 belong to the pre- recommendation and 56 cases belong to the post-recommendation phase of guideline implementation. Out of 222 patients, 102 belonged to the clean class and 120 belonged to the clean- contaminated class. 21 cases got infected, giving a surgical site infection of 9.45%. Pathogens involved in infected cases and antibiotic susceptibility pattern are mentioned in respectively.

The cases were also evaluated for studying the prescription pattern of prophylactic antibiotic in different types of surgery. It included appendectomy [37 (16.66%)], head and neck surgery [31 (13.96%)], cholecystectomy [20 (9.0%)], laparoscopic cholecystectomy [27 (12.16%)], Exploratory laparotomy [22 (9.90%)], Trendelenburg ligation [11 (4.95%)] and hernia surgery, which includes inguinal hernia repair [22 (9.90%)], paraumbilical hernia [11 (4.95%)], incisional hernia [14 (6.30%)], hernioplasty [16 (7.20%)] and herniorrhaphy [11 (4.95%)]. The final guideline for prophylactic antibiotic use is given in 25(11.26%) cases received a single preoperative dose, in 16 (7.20%) cases, the antibiotic was stopped within 24 hours of surgery and in 181 (81.53%) cases, the antibiotic was continued for 5-7 days after surgery. In all cases, prophylactic antibiotic was administered through the intravenous route only. Prophylactic antibiotics were administered at the time of induction in the operation theatre, in all the cases. Average duration of hospital stay was found to be 16.34 days in the infected group (21 cases) and 7.43 days in the non-infected group (201 cases).

A total of 56 cases were studied, of which 32 (57.14%) cases were of clean class and 24 (42.85%) were from clean-contaminated class. There was no incidence of surgical site infection. In clean surgeries like hernia, single dose amoxicillin clavulanic acid was the preferred option [12 (70.58%)]. Out of 6 Trendelenburg ligation surgeries, single dose amoxicillin-clavulanic acid [3 (50%)] and cefotaxime [3 (50%)] were the preferred options in conditions where ulcer was not present. Out of 9 cases of head and neck surgery, amoxicillin-clavulanic acid [6 (66.66%)] was the preferred option, followed by cefotaxime 3 (33.33%). Out of 24 clean contaminated surgeries, cefotaxime with metronidazole was the preferred option in 19 (79.16%) cases. Out of 56 cases, single preoperative dose were received by 22 (39.28%) cases, antibiotic was stopped within 24 hours of surgery in 32 (57.14%) cases and it was continued postoperatively for 5-7 days in 2 (3.57%) cases. All the 56 cases received prophylactic antibiotic through intravenous route. In all the 56 cases, antibiotic was administered at the time of induction in the operation theatre.

During the post- recommendation period, the choice of prophylactic antibiotics in different types of clean and clean- contaminated surgeries was consistent with the guidelines. Patients belonging to the pre and post- recommendation phase were of a similar group with respect to demographics, risk and type or procedures. Absence of SSIs during the post-recommendation period is not solely due to appropriate antibiotic prophylaxis usage. It is a well known fact that antibiotic prophylaxis is an adjunct to and not a substitute for good aseptic procedures in the surgical environment. Most of the SSIs arise from the patient's endogenous flora, which contaminates the wound by direct contact.^[20] Preparation of the patient for surgery was done meticulously, to decrease the microbiological burden on the patient's bowels, skin, respiratory tract or genital tract, depending on the procedure being performed. Shaving was done on the previous day of surgery. Patients were instructed to take bath and wear clean clothes on the day of surgery, as exogenous contamination of wounds is also important in the pathophysiology of SSI, particularly for clean surgical procedures, because surgeons do not incise mucous membranes or hollow viscera. Airborne bacteria originating from the patient or clothing or skin of the surgical team suffice to create SSI in these types of procedures, particularly when implants are being placed. Unless the blood sugar levels are well-controlled by Insulin administration, patients were not taken up for the surgical procedure.

All horizontal surfaces were cleaned every morning before any intervention; horizontal surfaces and all surgical items (e.g., tables, buckets) were cleaned between procedures. At the end of the working day, a complete cleaning of the operating theatre was performed. Once a week, a complete cleaning of the operating room area, including all annexes such as dressing rooms, technical rooms and cupboards were done. All members of the surgical team who would work on the operating field, were educated to scrub arms and hands with chlorhexidine, iodophors or hexachlorophene for at least 5 minutes before the first procedure of the day and for 2-5 minutes between subsequent procedures. Sterile gloves of good quality were used. Gloves were changed immediately after any accidental puncture. Postoperative dressing was done only in the dressing area. Members of the surgical team entering the operating room when the operation is about to begin or already underway, wear a mask and headgear that fully covers hair, sideburns and neckline. Scrub suits covered most bare skin to decrease shedding of microorganisms from uncovered skin. As individuals shed up to 10^9 epithelial cells per day, many of which carry bacteria, all personnel working in the operating room followed this practice. Any member of the surgical team who suffered from a skin lesion such as a boil, was refrained from working in the operating room. As a part of ISO (International Organization for Standardization) standardization for the hospital, healthcare professionals were adequately trained in maintaining good aseptic techniques. The duration of prophylactic antibiotic administration changed remarkably after the standardized recommendations. During the post- recommendation phase, prophylactic antibiotics were stopped after 24-48 hours of surgery. The development of guidelines helped in effective utilization, by preventing the overuse of prophylactic antibiotics.

A multifaceted educational intervention involving a team effort of healthcare professionals can have a significant effect on effective antibiotic utilization and reducing the incidence of surgical site infections. Local guidelines seem more likely to be accepted and followed than those developed nationally. Development of similar antibiotic guidelines is essential to prevent emergence of resistant pathogens, to rationalize the use of antibiotics in the most cost-effective manner and for preventing the occurrence of hospital- acquired infections. The present study has identified that the interventions done at a specific time by the teamwork of healthcare professionals, influence the prescribing appropriateness of surgical antibiotic prophylaxis. Several studies have highlighted the importance of a multidisciplinary approach in effective utilization of antibiotic prophylaxis.[\[21\]](#),[\[22\]](#) This study supports the team effort by surgeons, anesthetists, microbiologists, nurses and clinical pharmacists in the effective utilization of prophylactic antibiotics in hospital. The study also highlights the incidence of surgical site infections and prophylactic antibiotic usage in an Indian set-up.

CONCLUSION:

Multifaceted educational intervention involving a team effort of healthcare professionals can have a significant effect on effective antibiotic utilization and in reducing the incidence of surgical site infections. Local guidelines seem more likely to be accepted and followed than those developed nationally.

The following are the basic principles to be followed for prophylactic antibiotic usage:. The final decision regarding the benefits and risks of prophylaxis for an individual patient will depend on:The

patient's risk of surgical site infection. The potential severity of the consequences of surgical site infection. The effectiveness of prophylaxis in that operation

The consequences of prophylaxis for that patient (e.g., increased risk of colitis) Treatment policies should be based on local information about the epidemiology of drug-resistant bacteria. Implementation of a prophylaxis policy should not trigger an automatic change in treatment policy. Inappropriate prolongation of surgical prophylaxis can be reduced by use of specific order forms for surgical prophylaxis or recording of prophylaxis in single dose sections of existing drug prescription charts. Patients with a history of anaphylaxis or urticaria or rash, occurring immediately after penicillin therapy are at increased risk of immediate hypersensitivity to penicillins and should not receive prophylaxis with a beta-lactam antibiotic.

Policies for surgical prophylaxis that recommend beta-lactam antibiotics as first line agents, should also recommend an alternative for patients with allergy to penicillins or cephalosporins. Prophylaxis should be started preoperatively in most circumstances, ideally within 30 minutes of the induction of anesthesia. An additional dose of prophylactic agent is not indicated in adults, unless there is blood loss of up to 1500 ml during surgery or hemodilution of up to 15 ml/kg.

BIBLIOGRAPHY

1. Burke JP. Infection control - A problem for patient safety. *N Engl J Med* 2003;348:651-6.
2. Graham JC, Pedler SJ. Surgical antibiotic prophylaxis. *In* : Walker R, Edwards C, editors. *Clinical Pharmacy and Therapeutics*. 3rd ed. Churchill Livingstone: London; 2003. p. 569-81.
3. John WD, Salmaan K, Stephen WT. Antimicrobial prophylaxis in surgery. *In* : Joseph TD, Robert LL, Gary CY, editors. *Pharmacotherapy: A pathophysiologic approach*. 5th ed. McGraw Hill Publications: New York; 2002. p. 2111-22.
4. Sasse A, Mertens R, Sion JP, Ronveaux O, Bossens M, de Mol P, *et al* . Surgical prophylaxis in Belgian hospitals: Estimate of costs and potential savings. *J Antimicrob Chemother* 1998;41:267-72.
5. Nguyen D, Macleod WB, Phung DC, Cong QT, Nguy VH, van Nguyen H, *et al* . Incidence and predictors of surgical site infections in Vietnam. *Infect Control Hosp Epidemiol* 2001;22:485-92.
6. Sohn AH, Parvez FM, Vu T, Hai HH, Bich NN, Le Thu TA, *et al* . Prevalence of surgical site infections and patterns of antimicrobial use in a large tertiary care hospital in the Ho Chi Minh city, Vietnam. *Infect Control Hosp Epidemiol* 2002;23:382-7.
7. Saito T, Aoki Y, Ebara K, Hirai S, Kitamura Y, Kasaoka Y, *et al* . Surgical-site infection surveillance at a small-scale community hospital. *J Infect Chemother* 2005;11:204-6.

8. Haley RW, Culver DH, Morgan WM, White JW, Emori TG, Hooton TM. Identifying patients at high risk of surgical wound infection: A simple multivariate index of patient susceptibility and wound contamination. *Am J Epidemiol* 1985;121:206-15.
9. Olson MM, Lee JT Jr. Continuous 10 year wound infection surveillance. Results, advantages and unanswered questions. *Arch Surg* 1990;125:794-803.
10. Soletto L, Pirard M, Boelaert M, Peredo R, Vargas R, Gianella A, *et al* . Incidence of surgical-site infections and the validity of the National Nosocomial Infections Surveillance System risk index in a general surgical ward in Santa Cruz, Bolivia. *Infect Control Hosp Epidemiol* 2003;24:26-30.
11. Arias CA, Quintero G, Vanegas BE, Rico CL, Patino JF. Surveillance of surgical site infections: Decade of experience at a Colombian tertiary care center. *World J Surg* 2003;27:529-33.
12. Eriksen HM, Chugulu S, Kondo S, Lingaas E. Surgical-site infections at Kilimanjaro Christian Medical Center. *J Hosp Infect* 2003;55:14-20.
13. Jovanovic B, Mazic N, Mijoljevic V, Obrenovic J, Jovanovic S. Nosocomial infections in the intensive care units. *Vojnosanit Pregl* 2006;63:132-6.
14. Kasatpibal N, Jamulitrat S, Chongsuvivatwong V. Standardized incidence rates of surgical site infection: A multicenter study in Thailand. *Am J Infect Control* 2005;33:587-94.
15. Lilani SP, Jangale N, Chowdhary A, Daver GB. Surgical site infection in clean and clean-contaminated cases. *Indian J Med Microbiol* 2005;23:249-52.
16. Murthy R, Sengupta S, Maya N, Shivananda PG. Incidence of post operative wound infection and their antibiogram in a teaching and referral hospital. *Indian Med Sci* 1998;52:553-5.
17. del Giudice P, Blanc V, Durupt F, Bes M, Martinez JP, Counillon E, *et al* . Emergence of two populations of methicillin-resistant *Staphylococcus aureus* with distinct epidemiological, clinical and biological features from patients with community-acquired skin infections. *Br J Dermatol* 2006;154:118-24.
18. John MB, John EP, Thomas VB. Antibacterial prophylaxis in surgery. The surgeons guide to antimicrobial chemotherapy. Arnold Publications: London; 2000. p. 114-20.
19. Page CP, Bohnen JM, Fletcher JR, McManus AT, Solomkin JS, Wittmann DH. Antimicrobial prophylaxis for surgical wounds. Guidelines for clinical care. *Arch Surg* 1993;128:79-88.
20. Marie-Claude Roy. The operating room. *In* : Wenzel R, Brewer T, Butzler JP, editors. *A Guide to Infection Control in the Hospital*. International Society for Infectious Diseases: Boston; 2004. p. 114-9.

21. Matuschka PR, Cheadle WG, Burke JD, Garrison RN. A new standard of care: Administration of preoperative antibiotics in the operating room. *Am Surg* 1997;63:500-3.
22. Talon D, Mourey F, Touratier S, Marie O, Arlet G, Decazes JM, *et al* . Evaluation of current practices in surgical antimicrobial prophylaxis before and after implementation of local guidelines. *J Hosp Infect* 2001;49:193-8