

Assessment of the effect of periodontal treatment in patients with coronary artery disease: A pilot survey

Pavitra Rastogi, Rameshwari Singhal, Asmita Sethi, Avinash Agarwal¹,
V. K. Singh², Rishi Sethi³

Departments of Periodontics, Faculty of Dental Sciences, C.S.M. Medical University (Erstwhile K. G. Medical College Lucknow), ¹Medicine, ²Social and Preventive Medicine, ³Cardiology, C.S.M. Medical University, Lucknow, Uttar Pradesh, India

Address for correspondence: Dr. Pavitra Rastogi, Department of Periodontics, Faculty of Dental Sciences, C.S.M. Medical University, Lucknow, Uttar Pradesh, India.

E-mail: drpavitrarastogi@rediffmail.com

ABSTRACT

Background: Periodontitis is a chronic inflammatory condition believed to cause a low but long lasting systemic inflammatory reaction which in turn contributes to the development of atherosclerosis. Recent data suggests that around 40% cases of coronary artery disease remain unaccounted despite the identification of the classical risk factors. **Objectives:** To evaluate the efficacy of non surgical periodontal therapy on the levels of serum inflammatory markers in subjects with chronic periodontitis and known coronary artery disease. **Materials and Methods:** Twenty subjects with known coronary artery disease (CAD) were recruited from the Department of Cardiology, CSM Medical University, Lucknow, India, for this study. Periodontal disease was measured through the clinical parameters bleeding on probing (BOP) and probing depth (PD). All subjects received non surgical periodontal therapy that included oral hygiene instructions and meticulous scaling and root planing. Systemic levels of inflammatory markers such as high-sensitivity C reactive proteins (hsCRP), tumor necrosis factor- α (TNF- α), and white blood cell (WBC) counts were measured prior to and 1 month after periodontal therapy. **Results:** Subjects experienced significant reductions in bleeding on probing (BOP) and probing depth (PD), indicating improvement in overall periodontal health. In all subjects, high-sensitivity C reactive proteins (hsCRP), and WBC counts were reduced significantly,; however, tumor necrosis factor- α (TNF- α) levels showed no statistically significant reduction. **Conclusion:** In this study, periodontal treatment resulted in a significant decrease in bleeding on probing (BOP) and probing depth (PD), and this treatment lowered the serum inflammatory markers (hsCRP and WBC counts) in patients with coronary artery disease. This may result in a decreased risk for coronary artery disease in the periodontally treated patients.

Key words: Coronary artery disease, high-sensitivity C-reactive protein, periodontitis, tumor necrosis factor- α

INTRODUCTION

Non communicable diseases are taking an epidemic

form and will be a major cause of death in developing countries by the year 2020.^[1] Over 29.8 million people have coronary artery disease in India.^[1] Studies conducted in India show that every second person above 35yrs of age has periodontal pockets and 30% of total teeth extracted after 35yrs of age are due to periodontal disease.^[2] As a result of high prevalence of both coronary artery disease (CAD) and periodontal disease it has led to a hypothesis that these might be connected.^[3]

Framingham heart study has identified a list of classical

Access this article online	
Quick Response Code: 	Website: www.jcdronline.com
	DOI: 10.4103/0975-3583.95366

risk factors (hypertension, male gender, age, smoking, diabetes mellitus, and obesity). However, they were not sufficient to account for the etiology of this multifactorial pathological process. Although major improvements have been made in primary and secondary prevention of major risk factors of CAD, it continues to be highly prevalent. Around 40% cases of atherosclerosis cannot be attributed to the classical risk factors.

Due to these facts, there is an increased interest for considering chronic infections as a risk factor for atherosclerosis.^[4] As inflammatory mechanisms play a central role in mediating all phases of atherosclerosis and destructive periodontal disease, when left untreated they become a chronic inflammatory condition leading to an increase in local and systemic inflammatory mediators. Thus, perio systemic connection is a major current interest in periodontology.

Periodontal associated inflammatory process contributes to an increase in inflammatory mediators including tumor necrosis factor- α (TNF- α), C reactive proteins (CRP) and interleukin- 6 (IL- 6).^[5]

Epidemiological studies have associated plasma levels of IL-6 and TNF - α with cardio vascular risk factors and have associated IL- 6 levels with a risk of CAD.

Thus, the purpose of this interventional survey was to see the effect of periodontal treatment on levels of serum inflammatory markers CRP, TNF- α and white blood cell (WBC) counts.

MATERIALS AND METHODS

Twenty subjects were recruited from the Department of Cardiology, CSM Medical University, Lucknow, India, for the survey. Subjects with a chronic generalized periodontitis having coronary artery disease confirmed by clinical findings and ECG as interpreted by a cardiologist were included in the study.

Exclusion criteria included current smoking, smoking within last 6 months, diabetes and acute/chronic systemic diseases (e.g. influenza, rheumatoid arthritis, COPD, or kidney disease), antibiotics or anti inflammatory drug administration within the last 2 months, and pregnancy/lactation.

Clinical periodontal parameters

Probing depth was recorded at six sites per tooth (mesio

buccal, mid buccal, disto buccal, mesio lingual, mid lingual, and disto lingual).^[6] Probing depth was recorded to the nearest millimeter using UNC 15 probe.

The presence or absence of bleeding was recorded using Sulcular Bleeding Index (SBI, Muhlemann and Son 1971).^[7]

All subjects received oral prophylaxis and sublingual scaling and root planing. Oral hygiene instructions were given and the subjects were recalled 1 month after the last treatment visit.

Serum inflammatory markers

Serum high-sensitivity C reactive protein (hsCRP) was measured by using particle- enhanced turbidimetric assay (hsCRP, latex, COBAS, Integra 400 plus, ROCHE, Basel, Switzerland). Precipitate was determined turbidimetrically at 552 nm. The lower limit of assay was 0.1mg/ml.

TNF- α serum levels were determined by sensitive enzyme-linked immunosorbent assay (ELISA) using commercially available kit (R and D systems Inc, Minneapolis, MN, USA). The lower limit of detection was 0.06 pg/ml. The plates were read using an automated microplate reader (Bio Rad, Hercules, California, USA).

WBC count ($\times/10^9$) was obtained.

All the parameters (PD, BOP, hsCRP, TNF- α , WBC counts) were obtained at the baseline and 1 month post treatment.

Statistical analysis

The data collected were analyzed using non parametric Wilcoxon signed rank test. *P* values ≤ 0.05 were considered statistically significant.

RESULTS

There was a statistically significant reduction in BOP and PD post treatment. BOP was reduced by 28% and PD was reduced by 41% at the end of 1 month [Table 1].

hsCRP and WBC levels were significantly reduced post treatment. However, the levels of TNF α were not significantly reduced. The hsCRP, TNF- α levels and WBC counts reduced by 18, 2, and 14%, respectively [Table 2].

DISCUSSION

The present study was conducted with the purpose of detecting the role of non surgical periodontal therapy in

Table 1: Data on periodontal disease markers before and after treatment

Parameters	Baseline	Post treatment	Change
Bleeding on probing (%) (mean ± SD)	66.6 (± 1.4)	48.1 (± 5.7)	18.5
Probing depth(mm) (mean ± SD)	4.40 (± 0.80)	2.60 (± 0.78)	1.8

Table 2: Data on serum inflammatory markers before and after treatment

Inflammatory markers	Baseline	Post treatment	Change
hsCRP (mg/L) (mean ± SD)	2.54 (± 0.31)	2.08 (± 0.31)	0.46
TNF α (pg/ml) (mean ± SD)	1.79 (± 0.13)	1.59 (± 0.13)	0.20
WBCs (10 ⁹ /L) (mean ± SD)	8.04 (± 1.08)	6.90 (± 0.87)	1.14

subjects with coronary artery disease. The study clearly showed that there was a significant reduction in the levels of serum inflammatory markers (hsCRP and WBC counts) after periodontal therapy. The study is in agreement with the results of previous studies^[6-10] that stated a reduction in systemic inflammatory markers of inflammation associated with periodontitis post treatment.

In this study, non surgical therapy alone produced significant improvements in various periodontal parameters. BOP is an informative parameter to estimate the severity of gingival inflammation as well as the response to treatment, and it is a clinical indicator of disease progression and stability^[11]. BOP ≤ 20% of sites is associated with a lower risk for periodontal attachment loss. In the present study, BOP was significantly reduced 1 month after treatment, and the number of subjects with BOP >20% of sites at baseline was reduced by almost half after therapy. Similarly, all subjects experienced significant reductions in PD after the treatment.

In this study, 1month after mechanical therapy, the circulating levels of hsCRP and WBC counts significantly reduced in all subjects. CRP is a prognostic marker for future cardiovascular events.^[12] The release of bacteria and proinflammatory mediators such as bacterial endotoxins and cytokines in the bloodstream that causes the release of acute phase reactants (such as C reactive protein) leading to increased inflammatory activity in atherosclerotic lesions may represent the link between periodontal infection and CAD.

In addition, subjects with periodontitis present with higher counts of WBCs. WBC count, a crude marker of systemic inflammation associated with the prediction of future cardiovascular events^[13] also significantly decreased

for subjects with CAD. Individuals in our study showed a significant reduction of WBC after 1 month.

TNF-α has a role in apoptosis, bone resorption, matrix metalloproteinase (MMP) and IL-6 production. It has also been associated with increased risk of recurrent coronary events^[14]. However, the role of TNF-α remains disputed as several studies have not been able to recognize its association to cardiovascular events.^[15] Also, in our study there was no significant reduction in the levels of TNF-α. Large sample size is required to justify the association of TNF-α.

The observation that cardiovascular risk factors might be influenced by periodontitis may have important clinical consequences. First, as inflammation plays an important role in the pathophysiology of various conditions (metabolic syndrome, BP, vascular health).^[16] The association of mild chronic inflammation with future serious events in observational studies^[17] may be influenced by an underlying severe periodontal infection. Second, periodontitis may increase the risk of future cardiovascular events because of the pro atherogenic changes (increased cholesterol) and increased systolic blood pressure induced in affected individuals. Cigarette smoking represents the major influential factor with regard to the association between periodontal infections and systemic inflammation, and this preliminary investigation raises the hypothesis of a possible interaction of smoking, periodontal infection and systolic blood pressure on systemic health. Third, if periodontitis were the major inflammatory stimulus in at least some patients with periodontitis, severe periodontal infections may represent a major etiologic factor for atherosclerosis, metabolic syndrome, and their sequelae.

Still, the causal relationship between periodontitis and CAD cannot be established as the sample size was small and the role of other risk factors was not explored. On the basis of the data obtained, a large scale interventional study could be conducted in future.

CONCLUSION

- Poor oral health is detrimental to systemic health.
- Inflammatory markers are significantly associated with periodontitis.
- Periodontitis is a modifiable risk factor, which can be prevented and treated.

REFERENCES

1. Goyal A, Yusuf S. The burden of cardiovascular disease in the Indian subcontinent. *Indian J Med Res* 2006;124:235-44.
2. Sood M. A Study of Epidemiological Factors Influencing Periodontal

- Diseases in selected Areas of District Ludhiana, Punjab. Indian J Commun Med 2005;30:70-1.
3. Meurman J, Sanz M, Janket S. Oral Infection and Vascular Disease. Vascular Disease Prevention, 2007;4:260-7.
 4. Bretz W, Weyant RJ, Corby PM, Ren D, Weissfeld L, Kritchevsky SB. Systemic Inflammatory Markers, Periodontal Diseases, and Periodontal Infections in an Elderly Population. J Am Geriatrics Soc 2005;53:1532-7.
 5. Ide M, Jagdev D, Coward PY, Crook M, Barclay R, Wilson RF. The short term effects of treatment of chronic periodontitis on circulating levels of endotoxin, C-reactive protein, tumor necrosis factor- α and interleukin-6. J Periodontol 2004;75:420-8.
 6. Hussain Bokhari SA, Khan AA, Tatakis DN, Azhar M, Hanif M, Izhar M. Non-surgical periodontal therapy lowers serum inflammatory markers: a pilot study. J Periodontol 2009;80:1574-80.
 7. Muhleman HR, Son S. Gingival sulcus bleeding- a leading symptom in initial gingivitis. Helvetica Odontologica Acta 1971;33:107-13.
 8. D'Aiuto F, Parkar M, Andreou G, Suvan J, Brett PM, Ready D, *et al.* Periodontitis and Systemic Inflammation: Control of the Local Infection is Associated with a Reduction in Serum Inflammatory Markers. J Dent Res 2004;83:156-60.
 9. Mattila K, Vesanen M, Valtonen V, Nieminen M, Palosuo T, Rasi V, *et al.* Effect of treating periodontitis on C-reactive protein levels: a pilot study. BMC Infect Dis 2002;2:30-2.
 10. D'Aiuto F, Nibali L, Parkar M, Suvan J, Tonetti MS. Short-term effects of intensive periodontal therapy on serum inflammatory markers and cholesterol. J Dent Res 2005;84:269-73.
 11. Chaves ES, Wood RC, Jones AA, Newbold DA, Manwell MA, Kornman KS. Relationship of "bleeding on probing" and "gingival bleeding index" as clinical parameters of gingival inflammation. J Clin Periodontol 1993;20:139-43.
 12. Blake GJ, Ridker PM. Inflammatory bio-markers and cardiovascular risk prediction: Review. J Internal Medicine 2002;252:283-94
 13. Haim M, Boyko V, Goldbourt U, Battler A, Behar S. Predictive value of elevated white blood cell count in patients with preexisting coronary heart disease: The Bezafibrate Infarction Prevention Study. Arch Intern Med 2004;164:433-9.
 14. Ridker PM. Role of inflammatory biomarkers in prediction of coronary heart disease. Lancet 2001;358:946-8.
 15. Sukhija R, Fahdi I, Garza L, Fink L, Scott M, Aude W, *et al.* Inflammatory markers, angiographic severity of coronary artery disease, and patient outcome. Am J Cardiol 2007;99:879-84.
 16. Fernandez-Real JM, Ricart W. Insulin resistance and chronic cardiovascular inflammatory syndrome. Endocr Rev 2003;24:278-301.
 17. Ridker PM, Morrow DA. C-Reactive protein, inflammation, and coronary risk. Cardiol Clin 2003;21:315-25.

How to cite this article: Rastogi P, Singhal R, Sethi A, Agarwal A, Singh VK, Sethi R. Assessment of the effect of periodontal treatment in patients with coronary artery disease : A pilot survey. J Cardiovasc Dis Res 2012;3:124-7.
Source of Support: Nil, **Conflict of Interest:** None declared.

New features on the journal's website

Optimized content for mobile and hand-held devices

HTML pages have been optimized for mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed.

Click on **[Mobile Full text]** from Table of Contents page.

This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device.

Click on **[EPub]** from Table of Contents page.

There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops.

Links are available from Current Issue as well as Archives pages.

Click on  View as eBook